

Fred M. Katz

Sets and Their Sizes

New York, New York

Fred M. Katz (fred@katz.com)
Logic & Light, Inc.
New York, NY 10028
USA

Sets and Their Sizes

Mathematics Subject Classification (2001): 03E10, 54A25

©1981, 2001 Fred M. Katz.

All rights reserved. This work may not be translated or copied in whole or in part without the written permission of the publisher (Logic & Light, Inc. 400 East 84th Street, New York, NY 10028, USA), except for brief excerpts in connection with reviews or scholarly analysis.

The author hereby grants to M.I.T. permission to reproduce and distribute copies of this thesis document in whole or in part.

Preface

Cantor's theory of cardinality violates common sense. It says, for example, that all infinite sets of integers are the same size. This thesis criticizes the arguments for Cantor's theory and presents an alternative.

The alternative is based on a general theory, CS (for *Class Size*). CS consists of all sentences in the first order language with a subset predicate and a less-than predicate which are true in all interpretations of that language whose domain is a finite power set. Thus, CS says that *less than* is a linear ordering with highest and lowest members and that every set is larger than any of its proper subsets. Because the language of CS is so restricted, CS will have infinite interpretations. In particular, the notion of *one-one correspondence* cannot be expressed in this language, so Cantor's definition of similarity will not be in CS, even though it is true for all finite sets.

We show that CS is decidable but not finitely axiomatizable by characterizing the complete extensions of CS. CS has *finite completions*, which are true only in finite models and *infinite completions*, which are true only in infinite models. An infinite completion is determined by a set of *remainder principles*, which say, for each natural number, n , how many atoms remain when the universe is partitioned into n disjoint subsets of the same size.

We show that any infinite completion of CS has a model over the power set of the natural numbers which satisfies an additional axiom:

OUTPACING. If initial segments of A eventually become smaller than the corresponding initial segments of B , then A is smaller than B .

Models which satisfy OUTPACING seem to accord with common intuitions about set size. In particular, they agree with the ordering suggested by the notion of *asymptotic density*.

Acknowledgements

This dissertation has been made possible by Cantor — who invented set theory, Tarski — who discovered model theory, Abraham Robinson — who introduced model-completeness, and George Boolos — who made all of these subjects accessible to me. Prof. Boolos read several handwritten versions and as many typed versions of this thesis, with care and patience each time around. I owe

him thanks for finding (and sometimes correcting) my errors, for directing me to some useful literature, and simply for being interested. The reader should join me in thanking him for his help with style and organization, and for his insistence on filling in *mere details*. Nevertheless, whatever errors, gaps, or infelicities remain are solely the responsibility of the author.

Note

Except for some minor wording changes, typographical corrections, and omissions of some elementary (but very long) proofs, this is the text of my dissertation.

This dissertation was submitted in partial fulfillment of the requirements for the degree of Doctor of Philosophy at the Massachusetts Institute of Technology in April, 1981

It was certified and accepted by Prof. George Boolos, Thesis Supervisor and Chairperson of the Department of Linguistics and Philosophy.

Fred M. Katz
June, 2001

Contents

Preface	i
1 Introduction	1
1.1 The Problem	1
1.2 Cantor's argument	3
1.3 Cantor and the logicians	6
1.4 Aims and outline	7
2 The General Theory	9
2.1 The Theory CORE	9
2.2 Addition of Set Sizes	16
3 A Formal Theory of Class Size	19
3.1 CS - The Theory of Class Size	20
3.2 CA - Axioms for CS	22
3.3 Remarks on showing that CA axiomatizes CS	38
4 The Pure Theory of Class Sizes	45
4.1 Size models and PCS	45
4.2 Some models and theories	48
4.3 PCA is model complete	53
4.4 Remarks on showing that PCA axiomatizes PCS	61
5 Completeness of CA	63
5.1 Model completeness of CA	63
5.2 Prime models for CA	66
5.3 Summary	79
6 Sets of Natural Numbers	83
6.1 The outpacing principle	83
6.2 Models of CS and Outpacing	86
6.3 Size and density	94

Appendix	101
A Notation	101
B Model Theory	103
C Model Completeness	104
Bibliography	107
Index of Symbols	108
Index of Theories	110
Subject Index	112

1

Introduction

1.1 The Problem

This paper proposes a theory of set size which is based on intuitions, naive and otherwise. The theory goes beyond intuitions, as theories will, so it needs both justification and defense. I spend very little time justifying the theory; it is so clearly true that anyone who comes to the matter without prejudice will accept it. I spend a lot of time defending the theory because no one who comes to the matter comes without prejudice.

The prejudice stems from Cantor's theory of set size, which is as old as sets themselves and so widely held as to be worthy of the name *the standard theory*. Cantor's theory consists of just two principles:

ONE-ONE. Two sets are the same size just in case there is a one-one correspondence between them.

CANTOR[<]. A set, x , is smaller than a set, y , just in case x is the same size as some subset of y , but not the same size as y itself.

A **one-one correspondence** between two sets is a relation which pairs each member of either set with exactly one member of the other. For example, the upper-case letters of the alphabet can be paired with the lower-case letters:

ABCDEFGHIJKLMNOPQRSTUVWXYZ
abcdefghijklmnopqrstuvwxyz

So, the standard theory says, the set of upper-case letters is the same size as the set of lower-case letters. Fine and good.

The standard theory also says that the set of even numbers is the same size as the set of integers since these two sets can also be paired off one-to-one:

... $-n$... -3 -2 -1 0 1 2 3 ... n ...
... $-2n$... -6 -4 -2 0 2 4 6 ... $2n$...

Similarly, the standard theory says that the set of positive even integers is the same size as the set of prime numbers: pair the n -th prime with the n -th positive even number. In both of these cases, common sense chokes on the standard theory.

In the first case, common sense holds that the set of integers is larger than the set of even integers. The integers contain all of the even integers and then some. So it's just good common sense to believe there are more of the former than the latter. This is just to say that common sense seems to follow:

SUBSET. If one set properly includes another, then the first is larger than the second.

even into the infinite, where it comes up against the standard theory.

Common sense can make decisions without help from SUBSET. Though the set of primes is not contained in the set of even integers, it is still clear to common sense that the former is smaller than the latter. One out of every two integers is even, while the prime numbers are few and far between. No doubt, to use this reasoning, you need a little number theory in addition to common sense; but, given the number theory, it's the only conclusion common sense allows.

The theory proposed here accommodates these bits of common sense reasoning. It maintains SUBSET and a *few and far between* principle and much else besides. To state this theory, we use three two-place predicates: $<$, $\sim$, and $>$. If A and B name sets, then

- $\ulcorner A < B \urcorner$ is read as *A is smaller than B*,
- $\ulcorner A \sim B \urcorner$ is read as *A is the same size as B*, and
- $\ulcorner A > B \urcorner$ is read as *A is larger than B*.

Incidentally, we assume throughout this thesis that the following schemata are equivalent, item by item, to the readings of the three predicates given above, assuming that A is the set of α 's and B is the set of β 's:

- a.
 - There are fewer α 's than β 's.
 - There are just as many α 's than β 's.
 - There are more α 's than β 's.
- b.
 - The number of α 's is less than the number of β 's.
 - The number of α 's is the same as the number of β 's.
 - The number of α 's is greater than the number of β 's.
- c.
 - The size of A is smaller than the size of B .
 - A and B are (or, *have*) the same size.
 - The size of A is larger than the size of B .

Regarding this last group, we emphasize that we are not arguing that there *really are* such things as set sizes, nor that there *are not really* such things. Statements about *sizes* can be translated in familiar and long-winded ways into statements about sets, though we will not bother to do so.

I have identified the standard theory, Cantor's, with two principles about set size. The term *size*, however, is rarely used in connection with Cantor's theory; so it might be wondered whether the standard theory is really so standard. In stating ONE-ONE and CANTOR[<], Cantor used the terms *power* and *cardinal number* rather than *size*. In the literature, the term *cardinal number* (sometimes just *number*) is used most frequently. If someone introduces *cardinal number* as a defined predicate or as part of a contextual definition (e.g. "We say that two sets have the *same cardinal number* just in case . . . "), there is no point in discussing whether that person is right about *size*.

Though Cantor's theory is usually taken as a theory of set size, it can also be taken as *just* a theory of one-one correspondences. More specifically, saying that two sets are *similar* iff they are in one-one correspondence can either be taken as a claim about size or be regarded as a mere definition. Whether or not *similarity* is coextensive with *being the same size*, the definition is worth making. The relation picked out is well studied and well worth the study. The technical brilliance of the theory attests to this: it has given us the transfinite hierarchy, the continuum problem, and much else. In addition, the theory has consequences which do not *prima facie* seem to have anything to do with size or similarity: the existence of transcendental numbers comes to mind. All of this is to say that the interest in one-one correspondence has not been sustained solely by its identification with the notion of size. Hence, denying that they are the same does not endanger the theory of one-one correspondences, *per se*.

But most mathematicians and philosophers do not use *cardinal number* as a mere abbreviation. They use the term in just the way that we use *size* and slide freely among (A), (B), and (C). This is true, in particular, of Cantor, who offered ONE-ONE as a theory; indeed, he offered an argument for this theory.

1.2 Cantor's argument

Cantor bases his argument for ONE-ONE on the idea that the size of a set, it's cardinal number, depends on neither the particular elements it contains nor on how those elements are arranged:

The cardinal number, $|M|$, of a set M (is the general concept which) arises from M when we make abstraction of the nature of its elements and of the order in which they are given. [Cantor, p.86]

But to say that the cardinal number of a set does not depend on certain things is not to say what the cardinal is. Neither does it insure that two sets have the same cardinal number just in case they are in one-one correspondence. To flesh out this notion of *double abstraction*, Cantor reduces it to a second abstraction operator, one which works on the elements of sets rather than the sets

themselves:

Every element, m , if we abstract from its nature becomes a *unit*.
[Cantor, p.86]

and so, concludes Cantor:

The cardinal number, $|M|$, is a set composed of units (which has existence in our minds as an intellectual image or projection of M .)
[Cantor, p.86]

According to Jourdain, Cantor

distinguished very sharply between an aggregate and a cardinal number that belongs to it: “Is not an aggregate an object *outside* us, whereas its cardinal number is an abstract picture of it *in our* mind.”
[Cantor, p.80]

I have parenthesized the expressions above where Cantor describes cardinal numbers as mental entities. Nevertheless, I can only make sense of his arguments insofar as he treats cardinal numbers as sets: he refers to them as ‘definite aggregates’, supposes that they have elements, and employs mappings between cardinal numbers and other sets.

The following three statements seem to express Cantor’s intent:

$$(1.1) \quad |M| = \{ y \mid \exists x \in M \wedge y = \text{Abstract}(x) \}$$

$$(1.2) \quad \forall x \exists y (y = \text{Abstract}(x) \wedge \text{Unit}(y))$$

$$(1.3) \quad \forall M \forall y (y \in |M| \rightarrow \text{Unit}(y))$$

Abstract (x) is to be read as *the result of abstracting from the element x* , *Unit* (x) as *x is a unit*, and $|M|$ as *the cardinal number of M* .

So (1.1) gives a definition of cardinal number, in terms of the operation of abstraction, from which Cantor proves both ONE-ONEa and ONE-ONEb.

$$(ONE-ONEa) \quad M \sim N \rightarrow |M| = |N|$$

$$(ONE-ONEb) \quad |M| = |N| \rightarrow M \sim N$$

ONE-ONEa is true, says Cantor, because

the cardinal number $|M|$ remains unaltered if in the place of one or many or even all elements m of M other things are substituted.
[Cantor, p.80]

and so, if f is a one-one mapping from M onto N , then in replacing each element, m , of M with $f(m)$

M transforms into N without change of cardinal number. (p.88)

In its weakest form, the principle Cantor cites says that if a single element of M is replaced by an arbitrary element not in M then the cardinal number of the set will remain the same. That is,

$$(1.4) \quad (a \in M \wedge b \notin M) \\ \wedge N = \{x \mid (x \in M \wedge x \neq a) \vee x = b\} \\ \rightarrow |M| = |N|$$

The reasoning is clear: so far as the cardinal number of a set is concerned, one element is much the same as another. It is not the elements of a set, but only their abstractions, that enter into the cardinal number of a set. But abstractions of elements are just units; so one is much the same as another.

ONE-ONEb is true, Cantor says, because

... $|M|$ grows, so to speak, out of M in such a way that from every element m of M a special unit of $|M|$ arises. Thus we can say that $M \sim |M|$.

So, since a set is similar to its cardinal number, and similarity is an equivalence relation, two sets with same cardinal number are similar. Unless each element of a set abstracts to a 'special', i.e. distinct, unit, the correspondence from M to its cardinal number will be many-one and not one-one. A weak version of this principle is:

$$(1.5) \quad M = \{a, b\} \wedge a \neq b \\ \rightarrow |M| = \{\text{Abstract}(a), \text{Abstract}(b)\} \\ \wedge \text{Abstract}(a) \neq \text{Abstract}(b)$$

These two arguments do one another in. (1.4) says that replacing an element of a set with any element not in the set does not affect the cardinality. But, by the definition of $|M|$, (1.1), this means that

$$(1.6) \quad \forall x \forall y (\text{Abstract}(x) = \text{Abstract}(y))$$

For, consider an arbitrary pair of elements, a and b . Let $M = \{a\}$ and let $N = \{b\}$. So, the conditions of (1.4) are met and $|M| = |N|$. But $|M| = \{\text{Abstract}(a)\}$ and $|N| = \{\text{Abstract}(b)\}$, by (1.1). So $\text{Abstract}(a) = \text{Abstract}(b)$. Generalizing this argument yields (1.6).

So Cantor's argument for ONE-ONEa only works by assigning all non-empty sets the same, one-membered, cardinal number. But, this contradicts ONE-ONEb.

Conversely, the argument that a set is similar to its cardinal number relies on (1.5), which entails

$$(1.7) \quad \forall x \forall y (x \neq y \rightarrow \text{Abstract}(x) \neq \text{Abstract}(y))$$

assuming only that any two objects can constitute a set. But if the abstractions of any two elements are distinct, then no two sets have the same cardinal number as defined by (1.1), *contra* ONE-ONEa.

There is no way to repair Cantor's argument. Rather than leading to a justification of ONE-ONE, Cantor's definition of cardinal number is sufficient to refute the principle. The negation of (1.6) is:

$$(1.8) \quad \exists x \exists y (\text{Abstract}(x) \neq \text{Abstract}(y))$$

So one of (1.6) and (1.8) must be true. We have shown that (1.6) contradicts ONE-ONEb. Similarly, (1.8) contradicts ONE-ONEa; if a and b have distinct abstractions, then $\{a\}$ and $\{b\}$ have distinct cardinal numbers, $\{\text{Abstract}(a)\}$ and $\{\text{Abstract}(b)\}$, despite the fact that they are in one-one correspondence. So ONE-ONE is false whether (1.6) or its negation, (1.8), is true.

1.3 Cantor and the logicians

Though both Frege and Russell accepted Cantor's theory of cardinality, neither accepted Cantor's argument. Frege spends an entire chapter of the *Grundlagen* mocking mathematicians from Euclid to Schroder for defining numbers as sets of *units*. He neatly summarizes the difficulty with such views:

If we try to produce the number by putting together different distinct objects, the result is an agglomeration in which the objects remain still in possession of precisely those properties which serve to distinguish them from one another, and that is not the number. But if we try to do it in the other way, by putting together identicals, the result runs perpetually together into one and we never reach a plurality . . .

The word 'unit' is admirably adapted to conceal the difficulty . . . We start by calling the things to be numbered 'units' without detracting from their diversity; then subsequently the concept of putting together (or collecting, or uniting, or annexing, or whatever we choose to call it) transforms itself into arithmetical addition, while the concept word 'unit' changes unperceived into the proper name 'one'. [Frege, pp 50-51]

These misgivings about units do not prevent Frege from basing his definition of 'number' and his entire reduction of arithmetic on Cantor's notion of one-one correspondence. "This opinion", says Frege, "that numerical equality or identity must be defined in terms of one-one correlation, seems in recent years to have gained widespread acceptance among mathematicians" [Frege, pp. 73-74]. Frege cites Schroder, Kossak, and Cantor.

Russell displays similar caution about Cantor's argument [Russell, p.305] and similar enthusiasm for his theory (see the quote at the beginning of Chapter 2, for example.)

Of course, Frege and Russell *cleaned up* Cantor's presentation of the theory. Russell, for example, notes that Cantor's statement (1) is not a 'true definition' and

merely presupposes that every collection has some such property as that indicated — a property, that is to say, independent of its terms and their order; depending, we might feel tempted to add, only upon their number. [Russell, pp. 304-305]

So Russell, and similarly Frege, relied upon the principle of abstraction to obtain a 'formal definition' of cardinal numbers, in contrast to Cantor, who had "taken" number "to be a primitive idea" and had to rely on "the primitive proposition that every collection has a number." [Russell, p.305]

So, while some people regard Cantor's ONE-ONE as *just* a definition and others embrace it as a theory, the logicians have it both ways: adding ONE-ONE as a formal definition to set theory (or, as they would call it, logic) they have no obligation to defend it and can steer clear of peculiar arguments about *units*; at the same time, they can advance it as a great lesson for simple common sense.

The logicians' adoption of Cantor's theory of cardinality needs no great explanation: it *came with* set theory and, to a large extent, motivated set theory and determined its research problems. But there are two specific reasons that they should have seized upon ONE-ONE and CANTOR[<]. First, they both have the form of definitions, no matter how they are intended. So the notion of cardinality is *born reduced*.

Second, Cantor's theory clears the way for other reductions. Suppose, for example, you wish to reduce ordered pairs to sets. Well, you have to identify each ordered pair with a set and define the relevant properties and relations among ordered pairs in terms of properties and relations among sets. One of the relations that has to be maintained is *identity*; so each ordered pair must be identified with a distinct set. In addition, the relative sizes of sets of ordered pairs should be preserved under translation. But, if ONE-ONE is the correct theory of size, then this second condition follows from the first, since the existence of one-one correspondences will be preserved under a one-one mapping.

1.4 Aims and outline

It would be naive to suppose that people's faith in Cantor's theory would be shaken either by refuting specific arguments for ONE-ONE or by associating the acceptance of that theory with a discredited philosophy of mathematics. Such points may be interesting, but in the absence of an alternative theory of size they are less than convincing.

This dissertation presents such an alternative. Chapter 2 canvasses common sense intuitions for some basic principles about set size. Chapter 3 reorganizes those principles into a tidy set of axioms, offers an account of where the intuitions come from (viz. known facts about finite sets), and mines this source for additional principles. Chapters 4 and 5 prove that the theory so obtained is

complete, in the sense that it embraces all facts about finite sets of a certain kind (i.e. expressible in a particular language). Finally, Chapter 6 elaborates additional principles that concern only sets of natural numbers and demonstrates that these additional principles, together with the theory in Chapter 3, are satisfiable in the domain of sets of natural numbers.

2

The General Theory

The possibility that whole and part may have the same number of terms is, it must be confessed, shocking to common sense ... Common sense, therefore, is here in a very sorry plight; it must choose between the paradox of Zeno and the paradox of Cantor. I do not propose to help it, since I consider that, in the face of the proofs, it ought to commit suicide in despair. [Russell, p. 358]

Is common sense confused about set size, as Russell says, or is there a way of elaborating on common sense to get a plausible and reasonably adequate theory of cardinality? To be *plausible*, a theory should at least avoid principles and consequences which violate common sense. To be *reasonably adequate*, a theory has to go beyond bare intuitions: it should not rest with trivialities and it should answer as many questions about set size as possible, though it need not be complete. Plausibility and adequacy are conflicting demands: the first says that there should not be too many principles (no false ones, consistency), the second that there should not be too few principles.

In the Introduction, I argued that a coherent theory of cardinality has to contain some principles that refer to the kinds of objects in sets, *pace* Cantor. In this chapter, however, I want to see how far we can go without such principles; i.e. how much can you say about *smaller than* without using predicates (other than *identity*) which relate the members of the sets being compared? I shall begin by stating a number of principles and explaining why they are included in the general theory.

2.1 The Theory CORE

First, there is the SUBSET principle:

SUBSET. If x is a proper subset of y , then x is smaller than y .

The reason for including SUBSET should be obvious. What has prompted the search for an alternative to the standard theory of cardinality is the conflict

between ONE-ONE and SUBSET. Now, it is often said that common sense supports both of these principles and that it is in doing so that common sense is confused. From this it is supposed to follow that common sense cannot be relied upon, so we should opt for ONE-ONE, with the technical attractions that it provides.

But, there's a difference in the way that common sense supports these two principles. There is no doubt that you can lead an unsuspecting person to agree to ONE-ONE by focusing their attention on forks and knives, husbands and wives, and so forth: i.e. finite sets. With carefully chosen examples, say the odds and the evens, you might even convince someone that ONE-ONE is true for infinite sets, too. Now, I do not think that such guile is needed to lead someone to agree to SUBSET, but that's not what my argument depends on. The argument hinges on a suggestion about how to resolve cases where mathematical intuitions seem to conflict. The suggestion is to see what happens with particular cases on which the principles conflict *before* you've lead someone to agree to either of the general statements.

So, if you want to find out what common sense really thinks about SUBSET and ONE-ONE, you would present people with pairs of infinite sets, where one was a proper subset of the other. I've actually tried this, in an unscientific way, and what I've gotten, by and large, is what I expected: support for SUBSET. (*By and large* because many people think that all infinite sets have the same size: *Infinity*.)

Naturally, I would not venture that this sort of technique, asking people, is any way to find out which of SUBSET and ONE-ONE is *true*. People's intuitions about mathematics are notoriously unreliable, not to mention inconstant. Of course, harping on this fact might engender some unwarranted skepticism about mathematics. What I am suggesting is that there might be a rational way of studying mathematical intuitions and that we should at least explore this possibility before proclaiming common sense to be hopelessly confused on mathematical matters.

So, SUBSET, all by itself seems to be a plausible alternative to Cantor's theory. though it surely is not enough. Given just this principle, it's possible that one set is smaller than another just in case it is a proper subset of the other. It's clear that we need additional principles. All of the following seem worthy (where $x < y$ is to be read as *x is smaller than y*, $x \sim y$ is to be read as *x is the same size as y*, $x > y$ as *x is larger than y*. The Appendix gives a full account of the notations used throughout this thesis.)

Theory 2.1.1. QUASI-LOGICAL

(ASYM ^{<})	$x < y \rightarrow \neg y < x$
(ASYM ^{>})	$x > y \rightarrow \neg y > x$
(TRANS ^{<})	$(x < y \wedge y < z) \rightarrow x < z$
(TRANS ^{>})	$(x > y \wedge y > z) \rightarrow x > z$
(INDISC [~])	$x \sim y \rightarrow \text{Indisc}(x, y)$
(REF [~])	$x \sim x$
(SYM [~])	$x \sim y \rightarrow y \sim x$
(TRANS [~])	$(x \sim y \wedge y \sim z) \rightarrow x \sim z$
(DEF ^{>})	$x > y \leftrightarrow y < x$

where $\text{Indisc}(x, y)$ abbreviates $\forall z((z < x \leftrightarrow z < y) \wedge (z > x \leftrightarrow z > y))$

We call the principles listed above **quasi-logical principles** because it is tempting to defend them as logical truths. Consider the first principle, for example, in unregimented English:

ASYM[<]. If x is smaller than y , then y is not smaller than x .

This sentence can be regarded as an instance of the schema:

ASYM^F. If x is F -er than y , then y is not F -er than x .

where F is to be replaced by an adjective from which comparatives can be formed, e.g. ‘tall’, ‘short’, ‘happy’, but not ‘unique’ or ‘brick’. It appears that every instance of this schema is true, so it could be maintained that each is true in virtue of its form, that each is a logical truth.

The other principles might be defended in the same way, though the schema for $\text{INDISC}^{\sim}$ would have to be restricted to triples of corresponding comparatives, for example: *is smaller than, is larger than, is the same size as*.

But using such observations to support these principles would be problematic for two reasons. First, it would require taking positions on many questions about logical form and grammatical form which would take us far afield and, possibly, antagonize first-order logicians. Second, there are some instances of the schemata that make for embarrassing counterexamples: ‘further east than’ (in a round world) and ‘earlier than’ (in, I’m told, a possible world).

So, it might be that casting the principles above as instances of the appropriate schema would only explain why they are part of common sense. What remains clear is that a theory of cardinality which openly denied any of these principles would be implausible: it would be ridiculed by common sense and mathematical sophisticates alike. I can just barely imagine presenting a theory which, for fear of inconsistency, withheld judgment on one or more of these principles. But to do so without good reason would be counterproductive. It seems that if a case could be made that these statements, taken together, are inconsistent with SUBSET, that would be good reason to say that there is no

reasonably adequate alternative to Cantor's theory. Since my goal is to counter such a conclusion, it seems that the proper strategy is to include such seemingly obvious truths and to show that the resulting theory is consistent. So the strategy here is not to adduce principles and argue for the truth of each. This would be impossible, given that the principles are logically contingent. Instead, our approach is to canonize what common sense holds to be true about cardinality and show that the result is consistent and reasonably adequate.

As long as we restrict ourselves to SUBSET and the quasi-logical principles, consistency is no problem. After all, what do the quasi-logical principles say? Only that *smaller than* is a partial ordering, the *larger than* is the converse partial ordering, that *the same size as* is an equivalence relation, and that sets of the same size are indiscernible under the partial orderings. So, if we are given any domain of sets, finite or infinite, we get a model for our theory by assigning to $<$ the relation of *being a proper subset of*, assigning to $>$ the relation of *properly including*, and assigning to $\sim$ the identity relation. Since common sense knows that different sets can be the same size, there must be some additional principles to be extracted from common sense.

We shall now consider some principles which cannot be regarded as quasi-logical.

First, there is the principle of **trichotomy**.

$$(TRICH) \quad x < y \vee x \sim y \vee x > y$$

which says that any two sets are comparable in size. While a theory of set size which excluded TRICH might escape ridicule, it would surely be regarded with suspicion. Indeed, if the principles of common sense were incompatible with TRICH, this would undoubtedly be used to discredit them.

Second, there is the **representation principle**.

$$(REP^<) \quad x < y \rightarrow \exists x'(x' \sim x \wedge x' \subset y)$$

which says that if a set, x , is smaller than another, y , then x is the same size as some proper subset of y . Now, this is a principle which common sense has no particular feelings about. Analogous statements about physical objects are neither intuitive nor very clearly true. For example, (1) does not stand a chance of being regarded as true:

- (1) If one table is smaller than another, then the first is the same size as some proper part of the second.

if 'part' is taken to mean 'leg or top or rim or ...'. Even if common sense can be persuaded to take particles and arbitrary fusions of such as parts of tables, no one should condemn its residual caution about (1). If $REP^<$ is true, then it seems to be an interesting and special fact about sets.

$REP^<$ was originally included in this theory for technical reasons; it makes it easier to reduce the set of axioms already presented and it provides a basis for several principles not yet presented. $REP^<$ may be open to doubt, but it is not

a principle that Cantorians could complain about, for it is entailed by Cantor's definition of $<$:

$$(\text{CANTOR}^<) \quad x < y \leftrightarrow \neg(x \sim y) \wedge \exists x'(x' \sim x \wedge x' \subset y)$$

If $\text{CANTOR}^<$ is regarded as a principle instead of a definition, then it is entailed by the principles we have already mentioned:

If $x < y$, then $\neg(x \sim y)$, by $\text{INDISC}^\sim$ and $\text{ASYM}^<$. By $\text{REP}^<$, some proper subset of y , say x' , must be the same size as x . But $x' < y$, by SUBSET : so $x < y'$, by $\text{INDISC}^\sim$.

Conversely, if $x' \sim x$ and $x' \subset y$, then $x' < y$, by SUBSET . So $x < y$, by $\text{INDISC}^\sim$.

There are more principles to come, but before proceeding, I'd like to take stock of what we already have. First, I want to reduce the principles mentioned above to a tidy set of axioms. Second, I want to estimate how far we've gone.

The entire set of principles already adopted are equivalent to the following, which will be referred to as *the core theory*.

Theorem 2.1.2. CORE, the core theory,

$$\begin{aligned} (\text{SUBSET}) \quad & x \subset y \rightarrow x < y \\ (\text{DEF}^\sim) \quad & x \sim y \leftrightarrow \text{Indisc}(x, y) \\ (\text{TRANS}^\sim) \quad & (x \sim y \wedge y \sim z) \rightarrow x \sim z \\ (\text{DEF}^>) \quad & x > y \leftrightarrow y < x \\ (\text{IRREF}^<) \quad & \neg(x < x) \\ (\text{TRICH}) \quad & x < y \vee x \sim y \vee x > y \end{aligned}$$

The only axiom in CORE that has not already been introduced is $\text{DEF}^\sim$, which is logically equivalent to the conjunction of $\text{INDISC}^\sim$ and its converse $\sim\text{INDISC}$:

$$\begin{aligned} (\text{INDISC}^\sim) \quad & x \sim y \rightarrow \text{Indisc}(x, y) \\ (\sim\text{INDISC}) \quad & \text{Indisc}(x, y) \rightarrow x \sim y \end{aligned}$$

$\sim\text{INDISC}$ says that if two sets fail to be the same size, then their being different in size is attributable to the existence of some set which is either smaller than one but not smaller than the other, or larger than one but not larger than the other.

Theorem 2.1.3. *Let $T = \text{QUASI}; \text{SUBSET}; \text{REP}^<$. Then $T \equiv \text{CORE}$.*

Proof.

$\vdash T \vdash \text{CORE}$. We only need to show that $\sim\text{INDISC}$ is entailed by T . Suppose $\neg(x \sim y)$. So $x < y$ or $y < x$, by TRICH . But $\neg(x < x)$ and $\neg(y < y)$, by $\text{IRREF}^<$.

$\neg \text{CORE} \vdash T$

- (i) $\text{TRANS}^<$. If $y < z$, there is a y' such that $y' \sim y$ and $y' \subset z$ by $\text{REP}^<$. If $x < y$, then $x < y'$ because $y' \sim y$, by $\text{DEF}^\sim$. So, there is an x' such that $x' \sim x$ and $x' \subset y'$. So $x' \subset z$ and, by SUBSET , $x' < z$. But then $x < z$ by $\text{DEF}^\sim$.
- (ii) $\text{ASYM}^<$. If $x < y$ and $y < x$, then $x < x$, by $\text{TRANS}^<$, *contra* $\text{IRREF}^<$.
- (iii) $\text{TRANS}^>$, $\text{ASYM}^>$, and $\text{IRREF}^>$ follow from the corresponding principles for $<$ and $\text{DEF}^>$.
- (iv) $\text{INDISC}^\sim$, $\text{SYM}^\sim$, $\text{TRANS}^\sim$, and $\text{REF}^\sim$ are logical consequences of $\text{DEF}^\sim$.

□

CORE is consistent. In fact, two kinds of models satisfy CORE.

Definition 2.1.4.

a. $\mathcal{A}$ is a **finite class model** with basis x iff

- (i) $\dot{\mathcal{A}} = \mathcal{P}(x)$, where x is finite.
- (ii) $\mathcal{A} \models a \subset b$ iff $a \subset b$
- (iii) $\mathcal{A} \models a < b$ iff $|a| < |b|$

b. $\mathcal{A}$ is a **finite set model** iff

- (i) $\dot{\mathcal{A}} = \{x \mid x \text{ is a finite subset of } Y\}$, for some infinite Y .
- (ii) $\mathcal{A} \models a \subset b$ iff $a \subset b$
- (iii) $\mathcal{A} \models a < b$ iff $|a| < |b|$

Models can be specified by stipulation the *smaller than* relation since *larger than* and *same size as* are defined in terms of *smaller than*.

Fact 2.1.5.

- a. If $\mathcal{A}$ is a finite class model, then $\mathcal{A} \models \text{CORE}$.
- b. If $\mathcal{A}$ is a finite set model, then $\mathcal{A} \models \text{CORE}$.

Proof. In both cases, the finite cardinalities determine a quasi-linear ordering of the sets in which any set is higher than any of its proper subsets. □

The normal ordering of finite sets is, in fact, the only one that satisfies CORE. By adding TRICH we have ruled out all non-standard interpretations of $<$.

Theorem 2.1.6. Suppose that $\mathcal{A}$ is a model such that

- a. If $x \in \dot{\mathcal{A}}$, then x is finite,
- b. If $x \in \dot{\mathcal{A}}$ and $y \subset x$, then $y \in \dot{\mathcal{A}}$, and
- c. $\mathcal{A} \models \text{CORE}$.

Then $\mathcal{A} \models a < b$ iff $|a| < |b|$

Proof. We shall prove (*) by induction on n :

(*) If $|a| = n$, then $\mathcal{A} \models (a \sim b)$ iff $|b| = n$

Suppose	$n = 0$	
then	$a = \emptyset$	
But if	$ b = 0$	
then	$\mathcal{A} \models (a \sim b)$	by REF $\sim$
And if	$ b \neq 0$	
then	$a \subset b$	
So	$\mathcal{A} \models (a < b)$	by SUBSET
So	$\mathcal{A} \not\models (a \sim b)$	by INDISC $\sim$

Now, suppose that (*) is true for all $i \leq n$:

If	$ a = b = n + 1$	
then	$\mathcal{A} \not\models (a \sim b)$, as follows:	
Suppose	$\mathcal{A} \models (a < b)$	
So	$\mathcal{A} \models (a' \sim a)$ for some $a' \subset b$	by REP $<$
But if	$\mathcal{A} \models a' \subset b$	
then	$ a' \leq n$	
So	$ a \leq n$	by (*), contra our hypothesis.
But if	$ a = n + 1$	
and	$\mathcal{A} \models (a \sim b)$	
then	$ b \geq n + 1$	by induction.
But if	$ b > n + 1$	
pick	$b' \subset b, b' \in A$,	
with	$ b' = n + 1$	by condition (b)
So	$\mathcal{A} \models (b' < b)$	by SUBSET
and	$\mathcal{A} \models (b' \sim a)$	
So	$\mathcal{A} \models (a < b)$	contra our supposition.
So	$ b = n + 1$	

□

2.2 Addition of Set Sizes

We shall now extend CORE to an account of addition of set sizes. Since the domains of our intended models contain only sets and not sizes of sets, we have to formulate our principles in terms of a three-place predicate true of triples of sets: $\text{Sum}(x, y, z)$ is to be read as *the size of z is the sum of the sizes of x and y* .

The following principles are sufficient for a theory of addition:

Theory 2.2.1. Addition

Functionality of addition

- (a) $\text{Sum}(x, y, z) \rightarrow (x \sim x' \leftrightarrow \text{Sum}(x', y, z))$
- (b) $\text{Sum}(x, y, z) \rightarrow (y \sim y' \leftrightarrow \text{Sum}(x, y', z))$
- (c) $\text{Sum}(x, y, z) \rightarrow (z \sim z' \leftrightarrow \text{Sum}(x, y, z'))$

Addition for disjoint sets

$$(\text{DISJ}^+) \quad x \cap y = \emptyset \rightarrow \text{Sum}(x, y, x \cup y)$$

Monotonicity of Addition

$$(\text{MONOT}) \quad \text{Sum}(x, y, z) \rightarrow x < z \vee x \sim z$$

FUNC^+ says that sets bear Sum relations to one another by virtue of their sizes alone. This condition must clearly be met if Sum is to be read as specified above.

DISJ^+ tries to say what function on sizes the Sum relation captures by fixing the function on paradigm cases: disjoint sets. But FUNC^+ and DISJ^+ leave open the possibility that addition is cyclic: suppose we begin with a finite class model whose basis has n elements and assign to Sum those triples $\langle x, y, z \rangle$ where

$$|z| = (|x| + |y|) \bmod (n + 1)$$

Both FUNC^+ and DISJ^+ will be satisfied, though the interpretation of Sum does not agree with the intended reading. MONOT rules out such interpretations.

Given an interpretation of ' j ' over a power set there is at most one way of interpreting Sum which satisfies ADDITION . We shall show this by proving that ADDITION and CORE entail DEF^+ :

$$(\text{DEF}^+) \quad \text{Sum}(x, y, z) \leftrightarrow \exists x' \exists y' (x \sim x' \wedge y \sim y' \wedge x' \cap y' = \emptyset \wedge x' \cup y' = z)$$

DEF^+ says that the extension of Sum is determined by the extension of j .

A model of CORE must satisfy an additional principle, $\text{DISJ}^\cup$, if **Sum** is to be interpreted in a way compatible with **ADDITION**,

$$(\text{DISJ}^\cup) \quad (x \sim x' \wedge y \sim y' \wedge x \cap y = \emptyset \wedge x' \cap y' = \emptyset) \rightarrow (x \cup y) \sim (x' \cup y')$$

(Note: The proofs in this chapter will use boolean principles freely, despite the fact that we have not yet introduced them.)

Theorem 2.2.2. $\text{CORE} + \text{ADDITION} \vdash \text{DISJ}^\cup$

Proof.

Suppose $(x \sim x' \wedge y \sim y' \wedge x \cap y = \emptyset \wedge x' \cap y' = \emptyset)$
 then $\text{Sum}(x, y, x \cup y)$
 and $\text{Sum}(x', y', x' \cup y')$ by DISJ^+
 So $\text{Sum}(x', y, x \cup y)$ by FUNC^+ (a)
 So $\text{Sum}(x', y', x \cup y)$ by FUNC^+ (b)
 So $x \cup y \sim x' \cup y'$ by FUNC^+ (c)

□

If the minimal conditions on addition are to be satisfied in a model of CORE, then **Sum** has to be definable by DEF^+ .

Theorem 2.2.3. $\text{CORE} + \text{ADDITION} \vdash \text{DEF}^+$

Proof. $(\Rightarrow)$

Suppose $\text{Sum}(x, y, z)$
 So $x < z \vee x \sim z$ by **MONOT**
 But if $x \sim z$,
 Let $x' = z$
 and $y' = \emptyset$;
 then $\text{Sum}(x', y', z)$ by DISJ^+
 So $\text{Sum}(x, y', z)$ by FUNC^+ (a)
 So $y' \sim y$ by FUNC^+ (b)
 And if $x < z$
 pick $x' \subset z$
 with $x' \sim x$ by $\text{REP}^<$.
 Let $y' = z - x'$
 But $y' \sim y$, as before.

($\Leftarrow$)

Suppose	$(x \sim x' \wedge y \sim y' \wedge x' \cap y' = \emptyset \wedge x' \cup y' = z)$	
then	$\text{Sum}(x', y', z)$	by DISJ ⁺
So	$\text{Sum}(x, y', z)$	by FUNC ⁺ (a)
So	$\text{Sum}(x, y, z)$	by FUNC ⁺ (b)

□

Theorem 2.2.4. *EXCORE, the extended core, is CORE $\cup$ ADDITION.*

Fact 2.2.5. *The following are consequences of EXCORE:*

$$\begin{aligned}
 x \cap y_1 = x \cap y_2 = \emptyset \wedge y_1 \sim y_2 &\rightarrow x \cup y_1 \sim x \cup y_2 \\
 x \cap y_1 = x \cap y_2 = \emptyset \wedge y_1 < y_2 &\rightarrow x \cup y_1 < x \cup y_2 \\
 x_1 \cap y_1 = x_1 \cap y_2 = \emptyset \wedge x_1 \sim x_2 \wedge y_1 < y_2 &\rightarrow x_1 \cup y_1 < x_2 \cup y_2 \\
 x \subset z \wedge y \subset z \wedge x < y &\rightarrow (z - y) < (z - x) \\
 x \subset z \wedge y \subset z \wedge x \sim y &\rightarrow (z - y) \sim (z - x) \\
 x < y &\rightarrow \exists y'(y' \sim y \wedge x \subset y') \\
 x \sim y &\rightarrow x - (x \cap y) \sim y - (x \cap y) \\
 x < y &\rightarrow x - (x \cap y) < y - (x \cap y)
 \end{aligned}$$

Proof. The proofs are elementary.

□

3

A Formal Theory of Class Size

In the preceding chapter, we searched for principles that accord with pre-Cantorian ideas about sizes of sets. We produced several such principles, constituting EXCORE, and found two kinds of interpretations which satisfied these principles. In one case, the domains of the interpretations were finite power sets. In the other case, the domains consisted of all finite subsets of a given infinite set. Insofar as both kinds of interpretation have quantifiers ranging over finite sets, we may say that they demonstrate that EXCORE is true when it is construed as being about finite sets.

Our goal is to show that this general theory of set size can be maintained for infinite sets as well as for finite sets. We shall show this by constructing a model for the general theory whose domain is the power set of the natural numbers. But the ability to construct such a model is interesting only to the extent that the general theory it satisfies is reasonably adequate. Suppose, for example, that we offered as a general theory of size the axioms of CORE other than $\text{REP}^<$. Call this theory T . So T just says that *smaller than* is a quasi-linear ordering which extends the partial ordering given by the proper subset relation. Since any partial ordering can be extended to a quasi-linear ordering, T has a model over $\mathcal{P}(\mathbb{N})$. But unless we have a guarantee that the model constructed will satisfy, say, $\text{DISJ}^\cup$, the existence of the model does not rule out the possibility that T is incompatible with $\text{DISJ}^\cup$. For a particular principle, ϕ , in this case $\text{DISJ}^\cup$, we may take one of three tacks: (1) add ϕ to T , obtaining T' , and show that T' has a model over $\mathcal{P}(\mathbb{N})$; (2) show that ϕ is inconsistent with T and argue that T is somehow more fundamental or more intuitive than ϕ ; (3) acquiesce in ignorance of whether T and ϕ are compatible and argue that *if* they are incompatible, then T should be maintained anyway.

Below, we deal with $\text{DISJ}^\cup$ as in (1), since $\text{DISJ}^\cup$ is in EXCORE. Cantor's principle ONE-ONE is dealt with as in case (2). It seems futile to try to rule out the need to resort to the third approach for any cases at all, but we can

reduce this need to the extent that we include in our general theory, T , as many plausible statements as possible.

We cannot construct T by taking *all* statements which are true for finite sets. Not only is ONE-ONE such a statement, but using the notion of *all statements true for finite sets* presupposes that we have some idea of the range of *all statements*. To avoid the problems involved in speaking of *all statements*, we might instead settle for *all statements in L* , where L is some judiciously chosen language. To avoid ONE-ONE, T must fall short of the full expressive power of the language of set theory.

Consider, now, the axioms in EXCORE. Other than size relations, these axioms involve only boolean operations and inclusion relations among sets. They do not use the notions *ordered pair*, *relation*, or *function*. In short, the only set theory implicit in these axioms is boolean algebra, or a sort of *Venn diagram set theory*. This is not to say the axioms do not apply to relations, functions, or other sets of ordered pairs, but only that they do not refer to these sorts of objects as such.

In the next section, we define a language just strong enough to express EXCORE. We then construct a theory by taking all statements *in that language* which are true over all finite power sets. By drawing statements only from this relatively weak language, we arrive at a theory which can be satisfied over infinite power sets. But, since we include in the language *all* statements of the language which are true over any finite power set, we know that no statement in that language can arise as something which ought to be true over infinite power sets but might be incompatible with our theory.

There remains the possibility that we could follow the same strategy with a more expressive language, though it would have to remain less expressive than the full language of set theory. In fact, such a language can be obtained by including a notion of the product of set sizes. This in turn opens the possibility of a succession of richer languages and a corresponding succession of stronger theories of size. At this point, the possible existence of any such hierarchy is sheer speculation; we mention it only to emphasize that no claim is made here that we have the strongest possible general theory of set size.

3.1 CS - The Theory of Class Size

The theories discussed in this paper will be formulated within first order predicate logic with identity. To specify the language in which a theory is expressed, then, we need only list the individual constants, predicates, and operation symbols of the language and stipulate the *rank*, or number of argument places, for each predicate and each operation symbol.

Definition 3.1.1.

- a. L_C , L_C , *the language of classes*, is the first order language with individual constants $\emptyset$ and I , the one-place predicate, *Atom*, the two-place predicate $\subset$, and the two place operation symbols, $-$, $\cap$, and $\cup$.

- b. $L_{<}$, *the language of size*, is the first order language with the one-place predicate *Unit*, the two-place predicates $<$ and $\subset$, and the three place predicate *Sum*.
- c. $L_{C<}$, *the language of class size*, is the first order language containing just the non-logical constants in L_C and $L_{<}$.

Following the strategy outlined above, we define the theory of class size in terms of interpretations of $L_{C<}$ over finite power sets.

Definition 3.1.2.

- a. If $L_C \subset L$, then $\mathcal{A}$ is a **standard interpretation** of L iff
 - (i) $\dot{\mathcal{A}} = \mathcal{P}(x)$, for some x , and
 - (ii) $\mathcal{A}$ assigns the usual interpretations to all constants of L_C :

$$\begin{aligned}\mathcal{A}(\mathbf{I}) &= x, \\ \mathcal{A}(\emptyset) &= \emptyset, \\ \mathcal{A} \models a \subset b &\text{ iff } a \subset b\end{aligned}$$

- b. If $\mathcal{A}$ is a standard interpretation and $\dot{\mathcal{A}} = \mathcal{P}(x)$, then x is the **basis** of $\mathcal{A}$, $\mathcal{B}(\mathcal{A})$.

Definition 3.1.3. $\mathcal{A}$ is a **standard finite interpretation** of $L_{C<}$ iff

- a. $\mathcal{A}$ is a standard interpretation of $L_{C<}$,
- b. $\mathcal{A}$ has a finite basis, and
- c.

$$\begin{aligned}\mathcal{A} \models a < b &\text{ iff } |a| < |b|, \\ \mathcal{A} \models a \sim b &\text{ iff } |a| = |b|, \text{ and} \\ \mathcal{A} \models \text{Unit}(a) &\text{ iff } |a| = 1\end{aligned}$$

Definition 3.1.4. *CS, the theory of class size*, is the set of all sentences of $L_{C<}$ which are true in all standard finite interpretations of $L_{C<}$.

By drawing only on principles which can be stated in $L_{C<}$ we have at least ruled out the most obvious danger of paradox. Since the notion of one-to-one correspondence cannot be expressed in this language, Cantor's principle ONE-ONE will not be included in the theory CS, even though it is true over any finite power set.

Since CS has arbitrarily large finite models, it has infinite models. It is not obvious that CS has *standard* infinite models, in which the universe is an infinite power set. In Chapter 6 we show that such models do exist.

The present chapter is devoted to getting a clearer picture of the theory CS. Section 2 develops a set of axioms, CA, for CS. Section 3 outlines the proof that CA does indeed axiomatize CS. This proof is presented in Chapter 5, after a slight detour in Chapter 4.

3.2 CA - Axioms for CS

Here we shall develop a set of axioms, CA, for the theory CS. This will be done in several stages.

3.2.1 BA- Axioms for atomic boolean algebra

We'll begin with the obvious. Since all of the universes of the interpretations mentioned in the definition of CS are power sets, they must be atomic boolean algebras and, so, must satisfy BA:

Definition 3.2.1. *BA, the theory of atomic boolean algebras, is the theory consisting of the following axioms:*

$$\begin{aligned}
 x \cup y &= y \cup x \\
 x \cap y &= y \cap x \\
 x \cup (y \cup z) &= (x \cup y) \cup z \\
 x \cap (y \cap z) &= (x \cap y) \cap z \\
 x \cap (y \cup z) &= (x \cap y) \cup (x \cap z) \\
 x \cup (y \cap z) &= (x \cup y) \cap (x \cup z) \\
 x \cap (\mathbf{I} - x) &= \emptyset \\
 x \cup (\mathbf{I} - x) &= \mathbf{I} \\
 x \subset y &\leftrightarrow (x \cup y) = y \wedge x \neq y \\
 \text{Atom}(x) &\leftrightarrow (\forall y)(y \subset x \rightarrow y = \emptyset) \\
 x \neq \emptyset &\rightarrow (\exists y)(\text{Atom}(y) \wedge (y \subset x \vee y = x))
 \end{aligned}$$

These axioms are adapted from [Monk, Def 9.3, p. 141 and Def. 9.28, p. 151].

BA is clearly not a complete axiomatization of CS, since BA does not involve any size notions. But BA does entail all the sentences in CS which do not themselves involve size notions. To show this we need to draw on some established facts about the complete extensions of BA (in the language L_C). The key idea here is that complete extensions of BA can be obtained either by stipulating the finite number of atoms in a model or by saying that there are infinitely many atoms.

Definition 3.2.2. *For $n \geq 1$,*

a. $ATLEAST_n$ is a sentence which says that there are at least n atoms:

$$\exists x_1 \dots \exists x_n (\text{Atom}(x_1) \wedge \dots \wedge \text{Atom}(x_n) \wedge \{x_i \neq x_j \mid 0 < i < j \leq n\})$$

b. $EXACTLY_n$ is a sentence which says that there are exactly n atoms:

$$ATLEAST_n \wedge \neg ATLEAST_{n+1}$$

- c. *INF* is a set of sentences which is satisfied in all and only infinite models of BA:

$$INF = \{ ATLEAST_n \mid n > 1 \}$$

- d. $BA_n = BA; EXACTLY_n$

- e. $BAI = BA \cup INF$

Fact 3.2.3.

- a. For $n \geq 1$, BA_n is categorical. [Monk, Cor 9.32, p.152]
- b. For $n \geq 1$, BA_n is complete. (Immediate from above)
- c. For $n \geq 1$, an n -atom atomic boolean algebra is isomorphic to any standard finite interpretation of L_C with an n -element basis. [Monk, Prop. 9.30, p.151]
- d. BA is complete. [Monk, Theorem 21.24, p.360]
- e. BAI and the theories BA_n , $n \geq 1$, are the only complete, consistent extensions of BA.

Fact 3.2.4. If $BAI \vdash \phi$, then ϕ is true in some finite model of BA.

Proof. $BA \cup INF \vdash \phi$. By compactness, then, there is a k such that

$$BA \cup \{ ATLEAST_n \mid 1 \leq n \leq k \} \vdash \phi$$

So ϕ is true in any atomic boolean algebra with more than k atoms. $\square$

Theorem 3.2.5. If $CS \vdash \phi$, and $\phi \in L_C$, then $BA \vdash \phi$.

Proof. If $BA \not\vdash \phi$, then $\neg\phi$ is true in some atomic boolean algebra, $\mathcal{A}$.

If $\mathcal{A}$ is finite, then $\mathcal{A}$ is isomorphic to some standard finite interpretation $\mathcal{A}'$ of L_C . But, then $\neg\phi$ is true in $\mathcal{A}'$, so ϕ is not true in $\mathcal{A}'$ and $\phi \notin CS$.

If $\mathcal{A}$ is infinite, then $\neg\phi$ is consistent with BAI. But BAI is complete, so $BAI \vdash \neg\phi$. By 3.2.4, $\neg\phi$ is true in some finite model $\mathcal{A}$ of BA. Hence, $\neg\phi$ is true in some standard finite interpretation of L_C and, again, $\phi \notin CS$. $\square$

3.2.2 Size principles

Here, we just gather the principles presented above as EXCORE:

Theory 3.2.6. *CORE SIZE consists of the following axioms:*

(SUBSET)	$x \subset y \rightarrow x < y$
(DEF ^{>})	$x > y \leftrightarrow y < x$
(DEF [~])	$x \sim y \leftrightarrow \text{Indisc}(x, y)$
(IRREF ^{<})	$\neg(x < x)$
(TRICH)	$x < y \vee x \sim y \vee x > y$
(DEF ¹)	$\text{Unit}(x) \leftrightarrow \text{Atom}(x)$
(DEF ⁺)	$\text{Sum}(x, y, z) \leftrightarrow \exists x' \exists y' ($
	$x \sim x' \wedge y \sim y'$
	$\wedge x' \cap y' = \emptyset$
	$\wedge x' \cup y' = z)$
(DISJ ^U)	$(x \sim x' \wedge y \sim y'$
	$\wedge x' \cap y' = \emptyset$
	$\wedge x' \cup y' = z) \rightarrow x \cup y \sim x' \cup y'$

Combining the principles of boolean algebra and the size principles, we obtain our first serious attempt at a general theory of size:

Definition 3.2.7. *BASIC, the basic theory, is defined as:*

$$BA \cup \text{SIZE}$$

3.2.3 Division principles

BASIC is not a complete axiomatization of CS. In this section we shall exhibit an infinite number of principles which need to be added to BASIC in order to axiomatize CS. When we are done, we will have an effective set of sentences, **CA, the Class Size Axioms**, though we will not prove that CA $\equiv$ CS until Chapter 5.

To show that the new principles really do need to be added, we'll need some non-standard models of BASIC. These models will be similar in that (1) their universes will be subsets of $\mathcal{P}(\mathbb{N})$, (2) their atoms will be singletons in $\mathcal{P}(\mathbb{N})$, and (3) all boolean symbols will receive their usual interpretations. The models will, however, include different subsets of $\mathbb{N}$ and assign different size orderings to these sets.

In Chapter 6, these models of BASIC will reappear as submodels of various standard models of CS over $\mathcal{P}(\mathbb{N})$. So, in addition to the immediate purpose of establishing independence results, these models provide a glimpse of how sets of natural numbers are ordered by size.

Every standard finite interpretation, $\mathcal{A}$, of $L_{C<}$ satisfies exactly one of the

following:

$$\begin{aligned}
 (\text{EVEN}) \quad & \exists x \exists y (x \sim y \wedge x \cap y = \emptyset \wedge x \cup y = \mathbf{I}) \\
 (\text{ODD}) \quad & \exists x \exists y \exists z (x \sim y \wedge x \cap y = x \cap z = y \cap z = \emptyset \\
 & \wedge \text{Atom}(z) \wedge x \cup y \cup z = \mathbf{I})
 \end{aligned}$$

$\mathcal{A} \models \text{EVEN}$ if $|\mathcal{B}(\mathcal{A})|$ is even and $\mathcal{A} \models \text{ODD}$ if $|\mathcal{B}(\mathcal{A})|$ is odd. But $\text{BASIC} \not\models (\text{EVEN} \vee \text{ODD})$. Consider the model $\mathcal{F}$ whose universe consists of all only the finite and cofinite subsets of $\mathbb{N}$, where, for $a, b \in \mathcal{F}$:

$$\begin{aligned}
 \mathcal{F} \models (\mathbf{a} < \mathbf{b}) \text{ iff } & a \text{ and } b \text{ are both finite and } |a| < |b| \\
 & \text{or } a \text{ and } b \text{ are both cofinite and } |\mathbb{N} - a| > |\mathbb{N} - b| \\
 & \text{or } a \text{ is finite and } b \text{ is infinite.}
 \end{aligned}$$

$\mathcal{F}$ is a model of BASIC. But neither EVEN nor ODD is true in $\mathcal{F}$, for any two sets that are the same size are either both finite, in which case their union is also finite, or both cofinite, in which case they cannot be disjoint.

So $\text{EVEN} \vee \text{ODD}$ is in CS, but not entailed by BASIC. As you might suspect, this is just the tip of the iceberg of principles missing from an axiomatization of CS. Informally, we can extend $\mathcal{F}$ to a model that satisfies EVEN by including the set of even numbers and the set of odd numbers and making them the same size. To round out the result to a model of BASIC, we also need to include all sets which are *near* the set of evens or the set of odds, i.e. those that differ from the evens or odds by a finite set. With these additions made, the new model will be closed under boolean operations and will satisfy BA. There is a (unique) way of ordering these added sets by *smaller than* that will satisfy BASIC: rank them according to the size and direction of their finite difference from the odds or the evens. So, we can construct an infinite model of BASIC ; $(\text{EVEN} \vee \text{ODD})$.

But this model will still not satisfy CS, as we can see by generalizing the argument above. $(\text{EVEN} \vee \text{ODD})$ says that the universe is **roughly divisible** by two: EVEN says that the universe is divisible by two without remainder. ODD says that there is a remainder of a single atom. We can construct a similar statement that says the universe is roughly divisible by three — with remainder 0, 1, or 2. As with $(\text{EVEN} \vee \text{ODD})$, this statement is in CS but is satisfied by neither our original model nor the model as amended. Again, we can extend the model and again we can produce a statement of CS which is false in the resulting model.

We now formalize this line of reasoning.

Definition 3.2.8. *If $0 \leq m < n$, then*

a. $MOD_{n,m}$ is the sentence

$$\begin{aligned} \exists x_1 \dots \exists x_n \exists y_1 \dots \exists y_m (& \\ & x_1 \sim x_2 \sim \dots \sim x_n \\ & \wedge \text{Atom}(y_1) \wedge \dots \wedge \text{Atom}(y_m) \\ & \wedge \bigwedge_{1 \leq i < j \leq n} (x_i \cap x_j = \emptyset) \\ & \wedge \bigwedge_{1 \leq i < j \leq m} (y_i \cap y_j = \emptyset) \\ & \wedge (x_1 \cup \dots \cup x_n) \cap (y_1 \cup \dots \cup y_m) = \emptyset \\ & \wedge (x_1 \cup \dots \cup x_n) \cup (y_1 \cup \dots \cup y_m) = \mathbf{I} \end{aligned}$$

$MOD_{n,m}$ says that the universe is divisible into n sets of the same size with m atoms remaining.

b. DIV_n is the sentence

$$MOD_{n,0} \vee \dots \vee MOD_{n,n-1}$$

Fact 3.2.9. If $0 \leq m < n$ and $\mathcal{A}$ is a standard finite interpretation of $L_{C<}$, then

- (a) $\mathcal{A} \models MOD_{n,m}$ iff $|B(\mathcal{A})| \equiv m \pmod{n}$
- (b) $\mathcal{A} \models DIV_n$
- (c) $CS \vdash DIV_n$

Theorem 3.2.10. $BASIC \not\models CS$

Proof. If $n > 1$, $BASIC \not\models DIV_n$. The model, $\mathcal{F}$ defined above satisfies $BASIC$ but not DIV_n , for any n finite sets have a finite union and any two cofinite sets overlap. So, $BASIC \not\models CS$, by 3.2.9c. $\square$

We could consider adding all DIV_n sentences to $BASIC$ in the hope that this would yield a complete set of axioms for CS . We did consider this, but it does not work. To demonstrate this, we need some independence results for sets of DIV_n sentences.

Definition 3.2.11. If J is a set of natural numbers, then

- a. $DIV_J = \{ DIV_n \mid n \in J \}$
- b. $BDIV_J = BASIC \cup DIV_J$
- c. $BDIV_j = BDIV_{\{j\}}$

Our independence results will be obtained by constructing models of BASIC which satisfy specific sets of DIV sentences. To build such models from subsets of $\mathbb{N}$, we shall include sets which can be regarded as fractional portions of $\mathbb{N}$.

Definition 3.2.12. For $n \geq 0$,

- a. x is an ***n-congruence class*** iff $x = [n\mathbf{k} + m]$ for some m , where $0 \leq m < n$.
- b. x is an ***n-quasi-congruence class*** iff x is the union of finitely many n -congruence classes.
- c. x is a ***congruence class*** iff x is an n -congruence class for some n .
- d. x is a ***quasi-congruence class*** iff x is an n -quasi-congruence class for some n .
- e. $QC_n = \{ x \mid x \text{ is an } n\text{-quasi-congruence class} \}$
- f. $QC = \bigcup_{n \geq 0} QC_n$

Examples.

- a. The set of evens, $[2\mathbf{n}]$, and the set of odds, $[2\mathbf{n} + 1]$, are 2-congruence classes.
- b. $[3\mathbf{k} + 2]$ is a 3-congruence class.
- c. $\mathbb{N}$ is a 1-congruence class.
- d. $\mathbb{N}$ is an n -quasi-congruence class for every $n > 0$:

$$\mathbb{N} = [n\mathbf{k} + 0] \cup \dots \cup [n\mathbf{k} + n - 1]$$

Fact 3.2.13.

- a. If $x \in QC_n$ and $y \in QC_n$, then $x \cup y \in QC_n$.
- b. If $x \in QC_n$, then $\mathbb{N} - x \in QC_n$.

Proof.

- a. Suppose $x = a_1 \cup \dots \cup a_k$ and $y = b_1 \cup \dots \cup b_j$. Then $x \cup y = a_1 \cup \dots \cup a_k \cup b_1 \cup \dots \cup b_j$.
- b. $\mathbb{N}$ itself is the union of n -congruence classes. If x is the union of m of these classes, then $\mathbb{N} - x$ is the union of the remaining $(n - m)$ n -congruence classes.

□

Definition 3.2.14. x is ***near*** y , $NEAR(x, y)$, iff $x - y$ and $y - x$ are both finite.

Fact 3.2.15. *x is near y iff there are finite sets, w_1 and w_2 such that $x = (y \cup w_1) - w_2$.*

Proof.

$\Rightarrow$ Let $w_1 = x - y$ and let $w_2 = y - x$.

$\Leftarrow$ Suppose $x = (y \cup w_1) - w_2$. Then $x - y \subseteq w_1$ and $y - x \subseteq w_2$, so $x - y$ and $y - x$ are finite.

□

Fact 3.2.16. *If $x_1 \subseteq x \subseteq x_2$, x_1 is near y , and x_2 is near y , then x is near y .*

Proof. Since $x \subseteq x_2$, $(x - y) \subseteq (x_2 - y)$. But $(x_2 - y)$ is finite, so $(x - y)$ is finite.

Since $x_1 \subseteq x$, $(y - x) \subseteq (y - x_1)$. But $(y - x_1)$ is finite, so $(y - x)$ is finite. □

Fact 3.2.17. *NEAR is an equivalence relation.*

Proof.

- a. x is near x , since $x - x = \emptyset$, which is finite.
- b. If x is near y , then y is near x . Immediate.
- c. Suppose x is near y and y is near z . Note that

$$(z - x) = ((z \cap y) - x) \cup ((z - y) - x)$$

But $(z - y) - x$ is finite because $(z - y)$ is finite and $((z \cap y) - x)$ is finite because $((z \cap y) - x) \subseteq (y - x)$, which is finite. So the union, $(z - x)$, is finite.

Similarly,

$$(x - z) = ((x \cap y) - z) \cup ((x - y) - z)$$

where $((x \cap y) - z) \subseteq (y - z)$ and $((x - y) - z) \subseteq (x - y)$. So $(x - z)$ is also finite.

Hence, x is near z .

□

Fact 3.2.18.

- a. If x_1 is near x_2 , then $x_1 \cup y$ is near $x_2 \cup y$.
- b. If x_1 is near x_2 and y_1 is near y_2 , then $x_1 \cup y_1$ is near $x_2 \cup y_2$.
- c. If x is near y , then $\mathbb{N} - x$ is near $\mathbb{N} - y$

Proof.

- a. Since x_1 is near x_2 , $x_1 - x_2$ and $x_2 - x_1$ are finite. But

$$(x_1 \cup y) - (x_2 \cup y) \subseteq (x_1 - x_2)$$

$$\text{and } (x_2 \cup y) - (x_1 \cup y) \subseteq (x_2 - x_1)$$

- b. By (a), $x_1 \cup y_1$ is near $x_2 \cup y_1$, which is near $x_2 \cup y_2$. So $x_1 \cup y_1$ is near $x_2 \cup y_2$, by transitivity.
- c. $(\mathbb{N} - x) - (\mathbb{N} - y) = y - x$ and $(\mathbb{N} - y) - (\mathbb{N} - x) = x - y$. So if x and y are near each other, so are their complements.

□

We can now define the domains of the models we will use to establish the independence results.

Definition 3.2.19.

- a. $Q_n = \{ y \mid y \text{ is near an } n\text{-quasi-congruence class} \}$
- b. $Q = \bigcup_{n>0} Q_n$

Examples.

- a. $Q_1 = \{ y \subseteq \mathbb{N} \mid y \text{ is finite or cofinite} \}$.
- b. $Q_2 = \{ y \subseteq \mathbb{N} \mid y \text{ is finite, cofinite, near } [2\mathbf{n}] \text{ or near } [2\mathbf{n} + 1] \}$.
- Q_2 is the domain of the model constructed above to satisfy DIV_2 .

Fact 3.2.20. *If A is a class of sets such that*

- a. $\bigcup A \in A$,
- b. *If $x \in A$, then $\bigcup A - x \in A$, and*
- c. *If $x \in A$ and $y \in A$, then $x \cup y \in A$,*

then A forms a boolean algebra under the usual set-theoretic operations, where I is interpreted as A . [Monk, Def. 9.1, p.141 and Corr 9.4, p.142]

Theorem 3.2.21. *If $n > 0$, then Q_n forms an atomic boolean algebra under the usual set-theoretic operations.*

Proof. Using Fact 3.2.20:

- a. $\bigcup \{Q_n\} = \mathbb{N}$, since $\mathbb{N} \in \{Q_n\}$ and if $x \in Q_n$, then $x \subseteq \mathbb{N}$.

b. If $x \in Q_n$, then

Suppose x is near y
 and $y \in QC_n$
 So $\mathbb{N} - y \in QC_n$ by 3.2.13b
 and $\mathbb{N} - x$ is near $\mathbb{N} - y$ by 3.2.18b
 So $\mathbb{N} - x \in Q_n$.

c.

Suppose $x \in Q_n$
 and $y \in Q_n$
 then x is near x'
 and $x' \in QC_n$ for some x'
 and y is near y'
 and $y' \in QC_n$ for some y'
 But then $x' \cup y' \in QC_n$ by 3.2.13a
 and $x \cup y$ is near $x' \cup y'$ by 3.2.18b
 So $x \cup y \in QC_n$

□

Thus, Q_n is a boolean algebra. Moreover, every singleton is in Q_n since all singletons are near $\emptyset$. So Q_n is an atomic boolean algebra.

We now define a size function on all sets which are near quasi-congruence classes. The sizes assigned to sets are ordered pairs. The first member is a rational between 0 and 1 which represents the density of the set. The second member is an integer which represents the finite (possibly negative) deviation of a set from *average* sets of the same density. First, we define the ordering and arithmetic for these sizes with the intention of inducing the size ordering and Sum relation for sets from the assignment of sizes to sets.

Definition 3.2.22.

a. A **size** is an ordered pair $\langle \rho, \delta \rangle$, where ρ is rational and δ is an integer.

b. If $\theta_1 = \langle \rho_1, \delta_1 \rangle$ and $\theta_2 = \langle \rho_2, \delta_2 \rangle$ are sizes, then

(i) $\theta_1 < \theta_2$ iff $\rho_1 < \rho_2$ or $(\rho_1 = \rho_2 \wedge \delta_1 < \delta_2)$.

(ii) $\theta_1 + \theta_2 = \langle \rho_1 + \rho_2, \delta_1 + \delta_2 \rangle$

Only some of these sizes will actually be assigned to sets. Specifically, a size will be assigned to a set only if $0 \leq \rho \leq 1$. Moreover, if $\rho = 0$, then $\delta \geq 0$ and if $\rho = 1$, then $\delta \leq 0$.

Our intention in assigning sizes to sets is as follows: Suppose x is near an n -quasi-congruence class, x' . So x' is the union of $k \leq n$ n -quasi-congruence classes. The set x' has density k/n and this is the value, ρ , assigned to x . The δ value assigned to x is the finite number of elements added to or removed from x' to obtain x .

The definitions and facts below formalize this intention and demonstrate that the assignment of sizes to sets is well-defined.

Fact 3.2.23.

- a. If $x \in QC$, $y \in QC$, and $x \neq y$, then x is not near y . (That is, no two quasi-congruence classes are near each other.)
- b. Any set is near at most one quasi-congruence class.

Proof.

- a. Let n be the least number such that

$$x \in QC_n \text{ and } y \in QC_n$$

So each is a disjoint union of n -congruence classes:

$$\begin{aligned} x &= x_1 \cup \dots \cup x_j \\ \text{and } y &= y_1 \cup \dots \cup y_k \end{aligned}$$

Suppose $a \in x - y$
 then $a \in x_i$ for some i .
 But $a \notin y_j$, for any j .
 So $x_i \neq y_j$, for any j
 So $x_i \subseteq x - y$
 So $x - y$ is infinite, since x_i is infinite

Similarly, if $a \in y - x$, then $y - x$ is infinite.

- b. If x were near two quasi-congruence classes, the two would have to be near each other, since NEAR is transitive. But this is impossible by (a).

□

Definition 3.2.24. If x is near a quasi-congruence class, then

- a. $C(x)$ is the quasi-congruence class near x .
- b. $\Delta_1(x) = x - C(x)$.
- c. $\Delta_2(x) = C(x) - x$.

$\Delta_1(x)$ and $\Delta_2(x)$ are finite and

$$x = (C(x) \cup \Delta_1(x)) - \Delta_2(x)$$

Definition 3.2.25. If $x \in QC$, then

- a. $\alpha(x)$ = the least n such that $x \in QC_n$.
- b. $\beta(x)$ = the unique k such that x is the disjoint union of k $\alpha(x)$ -congruence classes.

Examples.

- a. If $x = [2\mathbf{n} + 1]$, $\alpha(x) = 2$ and $\beta(x) = 1$.
- b. If $x = [4\mathbf{n} + 1] \cup [4\mathbf{n} + 2]$, $\alpha(x) = 4$ and $\beta(x) = 2$.
- c. If $x = [4\mathbf{n} + 1] \cup [4\mathbf{n} + 3]$, $\alpha(x) = 2$ and $\beta(x) = 1$, since $x = [2\mathbf{n} + 1]$.

Definition 3.2.26. If $x \in Q$, then

$$\begin{aligned}\rho(x) &= \frac{\beta(C(x))}{\alpha(C(x))} \\ \delta(x) &= |\Delta_1(x)| - |\Delta_2(x)| \\ \theta(x) &= \langle \rho(x), \delta(x) \rangle\end{aligned}$$

We can, at last, define the models to be used in our independence proof.

Definition 3.2.27. For $n > 0$, $\mathcal{Q}_n$ is the interpretation $\mathcal{Q}$ of $L_{C<}$ in which boolean symbols receive their usual interpretation and

$$\begin{aligned}\dot{\mathcal{Q}} &= \mathcal{Q}_n \\ \mathcal{Q} \models x < y &\text{ iff } \theta(x) < \theta(y) \\ \mathcal{Q} \models x \sim y &\text{ iff } \theta(x) = \theta(y) \\ \mathcal{Q} \models \text{Unit}(x) &\text{ iff } \theta(x) = \langle 0, 1 \rangle \\ \mathcal{Q} \models \text{Sum}(x, y, z) &\text{ iff } \theta(z) = \theta(x) + \theta(y)\end{aligned}$$

To show that the models $\mathcal{Q}_n$ satisfy BASIC, we will need the following facts about congruence classes.

Fact 3.2.28.

- a. If $x = [a\mathbf{n} + b]$ and $a_2 = ac$, then

$$x = \bigcup_{0 \leq i < c} [a_2\mathbf{n} + (ia + b)]$$

- b. If $x \in QC_n$ and $m = kn$, then $x \in QC_m$.

c. If $x \in QC$ and $y \in QC$, there is an n such that $x \in QC_n$ and $y \in QC_n$.

Proof.

a. If $k \in [a_2\mathbf{n} + (ia + b)]$ for some i , $0 \leq i < c$, then, for some n_1 ,

$$\begin{aligned} k &= a_2n_1 + ia + b \\ &= a(cn_1) + ia + b \\ &= a(cn_1 + i) + b \end{aligned}$$

So $k \in x$.

If $k \in x$, there is an n_1 such that $k = an_1 + b$. Let n_2 be the greatest n such that $a_2n \leq k$ and let $k' = k - a_2n_2$.

Since $k \equiv b \pmod{a}$

and $a_2n_2 \equiv 0 \pmod{a}$

then $k' \equiv b \pmod{a}$

So $k' = ia + b$, where $0 \leq i < c$

But $k = a_2n_2 + k' = a_2n_2 + ia + b$

So $k \in [a_2\mathbf{n} + (ia + b)]$

b. By (a), each n -congruence class is a disjoint union of m -congruence classes.

c. Suppose $x \in QC_{n_1}$ and $y \in QC_{n_2}$. Then, by (b), both x and y are in $QC_{n_1n_2}$.

□

Theorem 3.2.29. For any $n > 0$, $\mathcal{Q}_n \models \text{BASIC}$

Proof. By 3.2.21, $\mathcal{Q}_n$ is an atomic boolean algebra; so $\mathcal{Q}_n \models \text{BA}$. The $<$ -relation of $\mathcal{Q}_n$ is induced from the linear ordering of sizes; so it is a quasi-linear ordering and $\text{IRREF}^<$, TRICH , and $\text{DEF}^\sim$ are satisfied. As for the remaining axioms:

a. **SUBSET**: Suppose $x \subset y$. If $C(x) = C(y)$ then $\rho(x) = \rho(y)$ and $\Delta_1(x) \subseteq \Delta_1(y)$ and $\Delta_2(x) \subseteq \Delta_2(y)$, where at least one of these inclusions is proper. So $\delta(x) < \delta(y)$.

But if $C(x) \neq C(y)$, then $C(x) \subset C(y)$, so $\rho(x) < \rho(y)$.

In either case, $\theta(x) < \theta(y)$, so $\mathcal{Q}_n \models x < y$.

b. **REP**[<]: Suppose $\mathcal{Q}_n \models x < y$. So $\theta(x) = \langle k_1/n, \delta_1 \rangle$, $\theta(y) = \langle k_2/n, \delta_2 \rangle$, and either $k_1 < k_2$ or $\delta_1 < \delta_2$.

We want to find some $x' \subset y$ such that $\mathcal{Q}_n \models x \sim x'$.

If $k_1 = k_2 > 0$, then y must be infinite; so, x' can be obtained by removing $\delta_2 - \delta_1$ atoms from y .

If $k_1 = k_2 = 0$, then $0 \leq \delta_1 < \delta_2$; so, again, x' can be obtained by removing $\delta_2 - \delta_1$ atoms from y .

If $k_2 > k_1 > 0$, then let y_1 be the union of k_1 n -congruence classes contained in $C(y)$. So $y - y_1$ is infinite and $y_1 - y$ is finite. Let $y_2 = y_1 - (y_1 - y) = y_1 \cap y$. So $\theta(y_2) = \langle k_1/n, -\delta_3 \rangle$ where $\delta_3 = |y_1 - y|$. Finally, let $\delta_4 = \delta_3 + \delta_1$ and $x' = y_2 \cup y_3$, where $y_3 \subseteq y_1 - y$ with $|y_3| = \delta_4$.

If $k_1 = 0$, then x is finite. If $k_2 > 0$, then y is infinite, so there is no problem. If $k_2 = 0$, then y is finite, but has more members than x , since $\delta_1 < \delta_2$. So, let x' be any proper subset of y with δ_1 members.

c. $\text{DISJ}^\cup$: It is enough to show that if x and y are disjoint, then $\theta(x \cup y) = \theta(x) + \theta(y)$. We need the following three facts:

(i) $C(x \cup y) = C(x) \cup C(y)$. (See Fact 3.2.18b.)

(ii) $\Delta_1(x \cup y) = (\Delta_1(x) \cup \Delta_1(y)) - (C(x) \cup C(y))$.

If $z \in x \cup y$ but $a \notin C(x \cup y)$, then $a \in \Delta_1(x)$ or $a \in \Delta_1(y)$; any element of $\Delta_1(x)$ is also in $\Delta_1(x \cup y)$ unless it is in $C(y)$; any element of $\Delta_1(y)$ is also in $\Delta_1(x \cup y)$ unless it is in $C(x)$.

(iii) $\Delta_2(x \cup y) = (\Delta_2(x) \cup \Delta_2(y)) - (\Delta_1(x) \cup \Delta_1(y))$.

Note that if $x \in Q$, $y \in Q$, and $x \cap y = \emptyset$, then $C(x) \cap C(y) = \emptyset$; otherwise, $C(x)$ and $C(y)$ have an infinite intersection.

Hence, if $a \in C(x) \cup C(y) - (x \cup y)$

then $a \in C(x) - x = \Delta_2(x)$

or $a \in C(y) - y = \Delta_2(y)$

And if $a \in \Delta_2(x)$, then $a \in \Delta_2(x \cup y)$

unless $a \in \Delta_1(y)$

And if $a \in \Delta_2(y)$, then $a \in \Delta_2(x \cup y)$

unless $a \in \Delta_1(x)$

.

From (ii) we obtain (iia):

$$(iia) \quad |\Delta_1(x \cup y)| = |\Delta_1(x) \cup \Delta_1(y)| \\ - |(\Delta_1(x) \cup \Delta_1(y)) \cap (C(x) \cup C(y))|$$

and from (iii) we obtain (iiia):

$$(iiia) \quad |\Delta_2(x \cup y)| = |\Delta_2(x) \cup \Delta_2(y)| \\ - |\Delta_1(x) \cup \Delta_1(y) \cap (\Delta_2(x) \cup \Delta_2(y))|$$

But $\Delta_1(x)$ and $\Delta_1(y)$ are disjoint, since x and y are disjoint. So:

$$(iv) \quad |\Delta_1(x) \cup \Delta_1(y)| = |\Delta_1(x)| + |\Delta_1(y)| = \delta_1(x) + \delta_1(y)$$

Since $\Delta_2(x)$ and $\Delta_2(y)$ are contained, respectively, in $C(x)$ and $C(y)$, which are disjoint, they are also disjoint. So:

$$(v) \quad |\Delta_2(x) \cup \Delta_2(y)| = \delta_2(x) + \delta_2(y)$$

So,

$$\begin{aligned} \delta_1(x \cup y) - \delta_2(x \cup y) &= (\delta_1(x) + \delta_1(y)) - (\delta_2(x) + \delta_2(y)) \\ &= (\delta_1(x) - \delta_2(x)) + (\delta_1(y) - \delta_2(y)) \\ &= \delta(x) + \delta(y) \end{aligned}$$

Since $\rho(x \cup y) = \rho(x) + \rho(y)$, by (i), we know that $\theta(x \cup y) = \theta(x) + \theta(y)$.

d. DEF⁺:

Suppose $\mathcal{Q}_n \models \text{Sum}(x, y, z)$
 So $\theta(z) = \theta(x) + \theta(y)$
 Clearly $\theta(x) \leq \theta(z)$
 Assume $\theta(x) = \theta(z)$
 then $\theta(y) = \langle 0, 0 \rangle$
 So $y = \emptyset$
 Let $x' = z$, $y' = \emptyset$ to satisfy DEF⁺
 Assume $\theta(x) < \theta(z)$
 then $\mathcal{Q}_n \models x' \sim x$
 and $x' \subset z$ for some x' since $\mathcal{Q}_n \models \text{REP}^<$
 Let $y' = z - x'$
 So $z = x' \uplus y'$
 Claim $\theta(y) = \theta(y')$ so $\mathcal{Q}_n \models y \sim y'$
 For $\theta(x') + \theta(y') = \theta(z)$ by DISJ^U
 But $\theta(x') = \theta(x)$
 So $\theta(y') = \theta(z) - \theta(x) = \theta(y)$

(Cancellation is valid for sizes because it is valid for rationals and integers.)

Conversely,

If $\theta(z) = \theta(x') + \theta(y')$
 and $\theta(x') = \theta(x)$
 and $\theta(y') = \theta(y)$
 then $\theta(z) = \theta(x) + \theta(y)$

□

Theorem 3.2.30. *For any $n > 0$, $\mathcal{Q}_n \models \text{DIV}_m$ iff $m \mid n$.*

Proof.

$\Rightarrow$ $\theta(\mathbb{N}) = \langle 1, 0 \rangle$. Hence, if m disjoint sets of the same size exhaust $\mathbb{N}$, they must each have size $\langle 1/m, 0 \rangle$. But if $x \in \mathcal{Q}_n$, then $\theta(x) = \langle a/n, b \rangle$, for integral a and b . So $b = 0$ and $a = n/m$.

$\Leftarrow$ For each i , $0 \leq i < n$, let $A_i = [n\mathbf{k} + i]$. So $\mathbb{N} = \bigcup_{0 \leq i < n} A_i$.

If $i \neq j$, then $A_i \cap A_j = \emptyset$ and $\mathcal{Q}_n \models (A_i \sim A_j)$ since $\theta(A_i) = \theta(A_j) = \langle 1/n, 0 \rangle$.

Letting $p = n/m$, group the n sets A_i into m collections with p members in each:

$$B_1, \dots, B_m$$

Letting $b_j = \bigcup B_j$ for $1 \leq j \leq m$, we have $b_j \in \mathcal{Q}_n$ and $\theta(b_j) = \langle p/n, 0 \rangle = \langle 1/m, 0 \rangle$.

Furthermore, $b_1 \cup \dots \cup b_m = \mathbb{N}$.

□

Definition 3.2.31. *If $J \neq \emptyset$ and J is finite, then the **least common multiple** of J , $\mu(J)$, is the least k which is divisible by every member of J .*

Remark. $\mu(J)$ always exists since the product of all members of J is divisible by each member of J . Usually, the product is greater than $\mu(J)$.

Corollary 3.2.32. *If J is finite, then*

- a. *If $\text{BDIV}_n \vdash \text{DIV}_m$, then $m \mid n$.*
- b. *If $\text{BDIV}_J \vdash \text{DIV}_m$, then $m \mid \mu(J)$.*
- c. *There are only finitely many m for which $\text{BDIV}_J \vdash \text{DIV}_m$.*

Proof. Based on Theorem 3.3.20:

- a. If $m \nmid n$, then $\mathcal{Q}_n \models \text{BDIV}_n; \neg \text{DIV}_m$
- b. $\mathcal{Q}_{\mu(J)} \models \text{BDIV}_J$ since it satisfies DIV_j for each $j \in J$. But, if $m \nmid \mu(J)$, then $\mathcal{Q}_{\mu(J)} \not\models \text{DIV}_m$.
- c. Immediate from (b), since only finitely many m divide $\mu(J)$.

□

We are now ready to show that $\text{BDIV}_n \not\vdash \text{CS}$ by finding a sentence in CS which entails infinitely many DIV_n sentences. Such sentences can be produced by generalizing the notion of divisibility to all sets instead of applying it only to the universe.

Definition 3.2.33.

a. If $0 \leq n$, then $\text{Times}_n(x, y)$ is the formula:

$$\exists x_0 \dots \exists x_n [x_0 = \emptyset \wedge x_n = y \wedge \bigwedge_{1 \leq i \leq n} \text{Sum}(x_{i-1}, x, x_i)]$$

$\text{Times}_n(x, y)$ says that y is the same size as the Sum of n sets, each the same size as x .

b. If $0 \leq m < n$, then $\text{Mod}_{n,m}(z)$ is the formula:

$$\exists x \exists y \exists v \exists w [\text{Times}_n(x, v) \wedge \text{Unit}(y) \wedge \text{Times}_m(y, w) \wedge \text{Sum}(v, w, z)]$$

$\text{Mod}_{n,m}(z)$ says that z can be partitioned in n sets of the same size and m atoms.

c. $\text{Div}_n(z)$ is the formula

$$\text{Mod}_{n,0}(z) \vee \dots \vee \text{Mod}_{n,n-1}(z)$$

d. ADIV_n is the sentence

$$\forall x \text{Div}_n(x)$$

Remark. We have taken this opportunity to formulate the divisibility predicates purely in terms of size predicates. Notice that in the presence of BASIC,

$$\begin{aligned} \text{MOD}_{n,m} &\equiv \text{Mod}_{n,m}(\mathbf{I}) \\ \text{and } \text{DIV}_n &\equiv \text{Div}_n(\mathbf{I}) \end{aligned}$$

Fact 3.2.34. $\text{CS} \vdash \text{ADIV}_n$, for every n .

Proof. Every set in every standard finite interpretation is a finite set, and all finite sets are roughly divisible by every n . $\square$

Fact 3.2.35. $\text{BASIC}; \text{ADIV}_n \vdash$:

- a. ADIV_{n^m} , for all m
- b. DIV_n , and
- c. DIV_{n^m} , for all m .

Proof.

- a. By induction on m : if $m = 1$, then $n^m = n$, so $\text{ADIV}_n \vdash \text{ADIV}_{n^m}$.
 If $T \vdash \text{ADIV}_{n^k}$, $\mathcal{A} \models T$, and $x \in \dot{\mathcal{A}}$, then x can be partitioned into n^k sets of the same size and i atoms, where $i < n^k$. Each non-atomic set in the partition can be further partitioned into n sets of the same size and j atoms, where $j < n$. Thus, we have partitioned x into $n^k n$ sets of the same size and $n^k j + i$ atoms. But $n^k n = n^{k+1}$ and, since $i < n^k$ and $j < n$, $n^k j + i < n^{k+1}$. Hence, $\mathcal{A} \models \text{ADIV}_{n^{k+1}}$.
- b. Obvious.
- c. Immediate from (a) and (b).

□

Theorem 3.2.36. $\text{BDIV}_{\mathbb{N}} \not\models \text{ADIV}_n$ for any $n \neq 1$.

Proof. If $\text{BDIV}_{\mathbb{N}} \vdash \text{ADIV}_n$, then by compactness there is a finite set J such that $\text{BDIV}_J \vdash \text{ADIV}_n$. But then $\text{BDIV}_J \vdash \text{DIV}_{n^k}$ for every k , by Fact 3.2.35c. But this contradicts Fact 3.2.32, which says that BDIV_J entails only finitely many DIV_n sentences. □

So, even if we add all of the DIV sentences to BASIC, we are left with a theory weaker than CS. Since this weakness has arisen in the case of ADIV sentences, it is reasonable to attempt an axiomatization of CS as follows:

Definition 3.2.37. $\text{CA} \equiv \text{BASIC} \cup \{ \text{ADIV}_n \mid n > 0 \}$

The remainder of this chapter and the next two are devoted to showing that CA is, indeed, a complete set of axioms for CS.

3.3 Remarks on showing that CA axiomatizes CS

We know that $\text{CS} \vdash \text{CA}$ and we want to show that $\text{CA} \equiv \text{CS}$, i.e. that $\text{CA} \vdash \text{CS}$. To do so, it will be sufficient to show that every consistent extension of CA is consistent with CS.

Fact 3.3.1.

- a. (*Lindenbaum's lemma*) Every consistent theory has a consistent, complete extension.
- b. If every consistent, complete extension of T_2 is consistent with T_1 , then $T_2 \vdash T_1$.

Proof.

- a. [Monk, Theorem 11.13, p.200]
- b. Suppose that $T_1 \vdash \phi$, $T_2 \not\vdash \phi$. Then $T = T_2; \neg\phi$ is consistent. T has a consistent, complete extension, T' , by Lindenbaum's lemma. Since $T_2 \subseteq T$, T' is also a consistent, complete extension of T_2 . But T' is not consistent with T_1 .

□

Definition 3.3.2. T' is a **completion** of T iff T' is a complete, consistent extension of T .

To prove that every completion of CA is consistent with CS, we define two kinds of completions of a theory.

Definition 3.3.3.

- a. T' is a **finite completion** of T iff T' is true in some finite model of T .
- b. T' is an **infinite completion** of T iff T' is true in some infinite model of T .

Fact 3.3.4. If T' is a completion of T and $BA \subseteq T$, then (T' is a finite completion of T iff T' is not an infinite completion of T .)

Proof.

$\Rightarrow$ Suppose $\mathcal{A}$ is a finite model of T' with n atoms. Since T' is complete, $T' \vdash \text{EXACTLY}_n$. So $T' \vdash \neg\text{ATLEAST}_{n+1}$ and has no infinite models.

$\Leftarrow$ $T' \vdash \text{ATLEAST}_n$ for every n , so T' has no finite models.

□

Fact 3.3.5.

- a. Every finite completion of CA is equivalent to $\text{CA}; \text{EXACTLY}_n$, for some n .
- b. Every finite completion of CA is consistent with CS.

Proof.

- a. $\text{CA} \vdash \text{BASIC}$ and, by Theorem 2.1.6, BASIC is categorical in every finite power.
- b. The model $\mathcal{A}$ of $\text{CA}; \text{EXACTLY}_n$ is a standard finite interpretation. So $\mathcal{A} \models \text{CS}$.

□

Definition 3.3.6.

- a. $CAI = CA \cup INF$
- b. $CSI = CS \cup INF$

So, to show that every completion of CA is consistent with CS, we may now concentrate on showing that every completion of CAI is consistent with CSI.

What, then, are the completions of CAI? Recall that CA entails DIV_n for every $n > 0$, where DIV_n is

$$MOD_{n,0} \vee \dots \vee MOD_{n,n-1}$$

Any completion, T , of CAI has to solve the disjunction DIV_n for each n ; T has to entail one of the disjuncts. *Remainder theories* specify, for each n , the number of atoms remaining when the universe is divided into n disjoint sets of the same size.

Definition 3.3.7. *Remainder functions and remainder theories*

- a. $f: \mathbb{N}^+ \rightarrow \mathbb{N}$ is a **remainder function** iff $0 \leq f(n) < n$ for all $n \in Dom(f)$. (Henceforth f ranges over remainder functions.)
- b. f is **total** iff $Dom(f) = \mathbb{N}^+$; otherwise f is **partial**.
- c. f is **finite** iff $Dom(f)$ is finite.
- d. n is a **solution** for f iff for any $i \in Dom(f)$, $n \equiv f(i) \pmod i$.
- e. f is **congruous** iff for any $i, j \in Dom(f)$, then $\gcd(i, j) \mid f(i) - f(j)$; otherwise, f is **incongruous**.
- f. The **remainder theory** specified by f , RT_f , is the set of sentences $\{ MOD_{n,m} \mid f(n) = m \}$.
- g. If T is a theory, $T_f = T \cup RT_f$.

Chapter 5 will show that if f is total, CAI_f is complete and that these are the only complete extensions of CAI. In this section, we will show that CAI_f is consistent just in case CSI_f is consistent.

Fact 3.3.8.

- a. If f is finite, then f has a solution iff f is congruous iff f has infinitely many solutions. [Griffin, Theorem 5-11, p.80]
- b. f is congruous iff every finite restriction of f is congruous.
- c. There are congruous f without any solutions. (Let $f(p) = p - 1$, for all primes p . Any solution would have to be larger than every prime.)

Theorem 3.3.9.

- a. If f is finite, then CS_f is consistent iff f is congruous.
- b. CS_f is consistent iff f is congruous.
- c. CSI_f is consistent iff f is congruous.

Proof.

- a. ($\Rightarrow$) Let $\phi = \bigwedge RT_f$. Since $CS; \phi$ is consistent, there is some n such that $\mathcal{A}_n \models \phi$. So n is a solution of f and, hence, f is congruous by 3.3.8a.
 ($\Leftarrow$) If f is congruous, f has a solution, n . So $\mathcal{A}_n \models CS; \phi$
- b. ($\Rightarrow$) For every finite restriction, g , of f , CS_g is consistent. By (a), each such g is congruous. Hence, f is congruous by 3.3.8b.
 ($\Leftarrow$) Every finite restriction, g , of f is congruous. So CS_g is consistent, by (a). By compactness, then, CS_f is consistent.
- c. ($\Rightarrow$) If CSI_f is consistent, so is CS_f . So, by (b), f is congruous.
 ($\Leftarrow$) By compactness, it is sufficient to show that every finite subtheory, T , of CSI_f is consistent. But, if T is such a theory, then

$$T \subseteq CS_g \cup \{ ATLEAST i \mid i < n \}$$

for some n and some finite restriction, g , of f . Since f is congruous, g is as well, by 3.3.8b. So g has arbitrarily large solutions and CS_g has finite models large enough to satisfy T . Hence, T is consistent.

□

We now want to prove a similar theorem for CA, our proposed axiomatization of CS. To do this, we must first establish that certain sentences are theorems of CA.

Lemma 3.3.10. *If $n \mid m$, $0 \leq q < n$, and $p \equiv q \pmod n$, then*

$$CA \vdash MOD_{m,p} \rightarrow MOD_{m,q}.$$

Proof. Suppose $\mathcal{A} \models MOD_{m,p}$, $k_1 = m/n$, and $p = k_2n + q$. So, $\mathcal{B}(\mathcal{A})$ can be partitioned into m sets of the same size

$$\begin{aligned} & b_{1,1} \dots b_{1,n} \\ & b_{2,1} \dots b_{2,n} \\ & \dots \\ & b_{k_1,1} \dots b_{k_1,n} \end{aligned}$$

and p atoms

$$\begin{aligned} & a_{1,1} \dots a_{1,n} \\ & a_{2,1} \dots a_{2,n} \\ & \dots \\ & a_{k_2,1} \dots a_{k_2,n} \\ & c_1 \dots c_q \end{aligned}$$

For $1 \leq i \leq n$, let

$$B_i = \bigcup_{1 \leq j \leq k_1} b_{j,n} \cup \bigcup_{1 \leq j \leq k_2} a_{j,n}$$

Since $\mathcal{A} \models \text{DISJ}^\cup$, $\mathcal{A} \models B_i \sim B_j$, for $1 \leq i, j \leq n$. Furthermore,

$$\mathcal{B}(\mathcal{A}) = \bigcup_{1 \leq i \leq n} B_i \cup (c_1 \dots c_q)$$

So, $\mathcal{A} \models \text{MOD}_{n,q}$. □

Lemma 3.3.11. *If $0 \leq p < q < m$, then*

$$CA \vdash \text{MOD}_{m,p} \rightarrow \neg \text{MOD}_{m,q}$$

Proof. Suppose $\mathcal{A} \models \text{MOD}_{m,p} \wedge \neg \text{MOD}_{m,q}$. Then:

$$\mathcal{A} \models x_1 \cup \dots \cup x_m \cup a_1 \cup \dots \cup a_p = I$$

$$\mathcal{A} \models y_1 \cup \dots \cup y_m \cup b_1 \cup \dots \cup b_q = I$$

where the a 's and b 's are atoms and the x 's (y 's) are disjoint sets of the same size in $\mathcal{A}$. Let

$$X = x_1 \cup \dots \cup x_m$$

$$Y = y_1 \cup \dots \cup y_m$$

$$A = a_1 \cup \dots \cup a_p$$

$$B = b_1 \cup \dots \cup b_p$$

$$B' = b_{p+1} \cup \dots \cup b_q$$

We claim that $y_1 < x_1$. For, if $y_1 \sim x_1$, then $X \cup A \sim Y \cup B$ and if $x_1 < y_1$, then $X \cup A < Y \cup B$; neither is possible since $Y \cup B \subset I = X \cup A$. So $y_i < x_i$ for $1 \leq i \leq m$. Since $\mathcal{A} \models \text{REP}^<$, there is a proper subset y'_i of x_i which is the same size as y_i .

$$\text{Let } z_i = x_i - y'_i$$

$$\text{and } Y' = y'_1 \cup \dots \cup y'_m$$

$$\text{So } Y' \cup Z = X = I - A$$

$$\text{and } Y' \cup B' = I - B.$$

But $A \sim B$, since each is the disjoint union of p atoms. Thus $(I - A) \sim (I - B)$, by $\text{RC}^\sim$, so $Y' \cup Z \sim Y' \cup B'$. But $Y' \sim Y$, since for each component of Y , there is a component of Y' of the same size. So $Z \sim B'$, by $\text{RC}^\sim$.

But Z must be larger than B' , for B' is the union of fewer than m atoms while Z is the union of m non-empty sets. So, the original supposition entails a contradiction. □

Theorem 3.3.12.

- a. If f is finite, then CA_f is consistent iff f is congruous.
- b. CA_f is consistent iff f is congruous.
- c. CAI_f is consistent iff f is congruous.

Proof.

- a. ($\Rightarrow$) Supposing that f is incongruous, there exist i, j , and k such that $k = \gcd(i, j)$ and $k \nmid (f(i) - f(j))$. We will show that

$$(*) \quad CA \vdash \neg(\text{MOD}_{i,f(i)} \wedge \text{MOD}_{j,f(j)})$$

from which it follows that CA_f is inconsistent.

Let p and q be such that

$$\begin{aligned} 0 &\leq p, q < k, \\ f(i) &\equiv p \pmod{k}, \text{ and} \\ f(j) &\equiv q \pmod{k} \end{aligned}$$

By lemma 3.3.10, we have

- (1) $CA \vdash \text{MOD}_{i,f(i)} \rightarrow \text{MOD}_{k,p}$, and
- (2) $CA \vdash \text{MOD}_{j,f(j)} \rightarrow \text{MOD}_{k,q}$

since $k \mid i$ and $k \mid j$. Since $k \nmid f(i) - f(j)$, $p \neq q$. So lemma 3.3.11 yields

$$(3) \quad CA \vdash \text{MOD}_{k,p} \rightarrow \neg \text{MOD}_{k,q}$$

From (1), (2), and (3) we may conclude (*).

($\Leftarrow$) Follows from 3.3.9a since $CA_f \subseteq CS_f$.

- b. See proof of 3.3.9b.

- c. See proof of 3.3.9c.

□

Corollary 3.3.13. CAI_f is consistent iff CSI_f is consistent.

Proof. Immediate from 3.3.9c and 3.3.12c.

□

It might help to review our strategy before presenting the difficult parts of the proof that $CA \equiv CS$. The main objective is (1), which follows from (2) by 3.3.1b.

$$(1) \quad CA \vdash CS$$

- (2) Every completion of CA is consistent with CS.

We already know that the finite completions of CA are consistent with CS (see 3.3.5c) and that if CAI_f is consistent, then CSI_f is also consistent (see 3.3.12). So (2) is a consequence of (3).

- (3) If T is a completion of CAI, then $T \equiv \text{CAI}_f$ for some total, congruous f .

To establish (3), it is sufficient to prove (4) because every completion of CAI entails CAI_f for some total f .

- (4) If f is total and congruous, then CAI_f is complete.

To prove (4), we invoke the prime model test: If T is model complete and T has a prime model, then T is complete (see Appendix, Fact C.3). So (4) follows from (5) and (6).

- (5) If f is total and congruous, then CAI_f has a prime model.

- (6) For any f , CAI_f is model complete.

Finally, since any extension of a model complete theory is also model complete (see Appendix, Fact C7a), we can infer (6) from (7).

- (7) CAI is model complete.

So (1) follows from (5) and (7).

The proof outlined here will be carried out in Chapter 5. But first, we consider a simpler theory, PSIZE, which deals only with sizes of sets and ignores boolean relations. Chapter 4 formulates PSIZE and establishes that it is model complete, a result we need for showing that CAI is model complete.

4

The Pure Theory of Class Sizes

CS is about sets; it makes claims about sets in terms of their boolean relations and their size relations. In this chapter, we identify a theory, PCS (the pure theory of class sizes), which is not about sets, but only about sizes of sets. *Sizes*, here, are equivalence classes of sets which have the same size. PCS is formulated in the language of size relations, $L_<$.

PCS is worth examining in its own right, for if number theory is the theory of cardinal numbers, then PCS is our version of number theory. But our main reason for introducing PCS is to aid the proof that CA is model complete. For this reason, we only give a sketchy treatment of PCS itself.

Section 1 defines PCS and develops a set of axioms, PCA, for it, as follows: for each model, $\mathcal{A}$, of BASIC, the **size model**, $\mathcal{S}_{\mathcal{A}}$, consists of equivalence classes drawn from $\dot{\mathcal{A}}$ under the *same size* relation; PCS is the set of statements true in $\mathcal{S}_{\mathcal{A}}$ for any standard finite model, $\mathcal{A}$. PCA consists of a theory, PSIZE, which holds in $\mathcal{S}_{\mathcal{A}}$ whenever $\mathcal{A} \models \text{BASIC}$ and a set of divisibility principles.

Using some results about model theory in section 2, Section 3 establishes that PCA is model complete, the main result of this chapter and the only result needed for subsequent proofs. This is done by reducing PCA to the theory Zgm, whose models are $\mathbb{Z}$ -groups taken modulo some specific element.

Finally, section 4 indicates how PCA could be shown to axiomatize PCS. This method is the same outlined in chapter 3 to show that $\text{CA} \equiv \text{CS}$.

4.1 Size models and PCS

Definition 4.1.1. *Suppose $\mathcal{A} \models \text{BASIC}$.*

- a. If x is a member of $\dot{\mathcal{A}}$, then $\sigma_{\mathcal{A}}[x]$ is the **size** of x in $\mathcal{A}$:*

$$\sigma_{\mathcal{A}}[x] = \{ y \mid \mathcal{A} \models (x \sim y) \}$$

b. $\mathcal{S}_{\mathcal{A}}$, the **size model** for $\mathcal{A}$, is the interpretation of $L_{<}$ whose domain is $\{\sigma_{\mathcal{A}}[x] \mid x \in \dot{\mathcal{A}}\}$ where

$$\begin{aligned} \mathcal{S}_{\mathcal{A}} \models \sigma_{\mathcal{A}}[x] \sim \sigma_{\mathcal{A}}[y] & \text{ iff } \mathcal{A} \models x \sim y \\ \mathcal{S}_{\mathcal{A}} \models \sigma_{\mathcal{A}}[x] < \sigma_{\mathcal{A}}[y] & \text{ iff } \mathcal{A} \models x < y \\ \mathcal{S}_{\mathcal{A}} \models \text{Sum}(\sigma_{\mathcal{A}}[x], \sigma_{\mathcal{A}}[y], \sigma_{\mathcal{A}}[z]) & \text{ iff } \mathcal{A} \models \text{Sum}(x, y, z), \text{ and} \\ \mathcal{S}_{\mathcal{A}} \models \text{Unit}(\sigma_{\mathcal{A}}[x]) & \text{ iff } \mathcal{A} \models \text{Atom}(x) \end{aligned}$$

c. The one-place operator $\tilde{x}$ is to be read as the complementary size of x :

$$\mathcal{S}_{\mathcal{A}} \models (y = \tilde{x}) \text{ iff } \mathcal{S}_{\mathcal{A}} \models \text{Sum}(x, y, \mathbf{I})$$

These interpretations are well defined because the predicates are satisfied by elements of $\dot{\mathcal{A}}$ in virtue of their sizes. For example, if $\mathcal{A} \models \text{Sum}(x, y, z)$ and $\mathcal{A} \models z \sim z'$, then $\mathcal{A} \models \text{Sum}(x, y, z')$.

Definition 4.1.2. *PCS, the pure theory of class sizes, consists of all sentences of $L_{<}$ which are true in the size model of every standard finite interpretation of $L_{<}$.*

Definition 4.1.3. *PSIZE, the theory of sizes, consists of the following axioms:*

Order axioms

$$\begin{array}{ll} (\text{IRREF}^{<}) & \neg(x < x) \\ (\text{TRANS}^{<}) & x < y \wedge y < z \rightarrow x < z \\ (\text{UNIQ}^{\sim}) & x \sim y \leftrightarrow x = y \\ (\text{MAX}) & x \leq \mathbf{I} \\ (\text{MIN}) & 0 \leq x \\ (\text{TRICH}) & x < y \vee x \sim y \vee y < x \end{array}$$

Unit axioms

$$\begin{aligned} \text{Unit}(x) & \leftrightarrow (y < x \leftrightarrow y = 0) \\ & \exists x \text{Unit}(x) \end{aligned}$$

Sum axioms

(IDENT)	$Sum(x, 0, x)$
(COMM)	$Sum(x, y, z) \leftrightarrow Sum(y, x, z)$
(MONOT)	$Sum(x_1, y, z_1) \wedge Sum(x_2, y, z_2) \rightarrow$ $(x_1 < x_2 \leftrightarrow z_1 < z_2)$
(ASSOC)	$Sum(x, y, w_1) \wedge Sum(w_1, z, w) \wedge Sum(y, z, w_2)$ $\rightarrow Sum(x, w_2, w)$
(EXIST ⁺)	$\exists z Sum(x, y_1, z) \wedge y_2 \leq y_1 \rightarrow \exists z Sum(x, y_2, z)$
(EXIST ⁻)	$x \leq z \rightarrow \exists y Sum(x, y, z)$
(COMP)	$Sum(x, \tilde{x}, \mathbf{I})$

Fact 4.1.4. *If $\mathcal{A} \models \text{BASIC}$, then $\mathcal{S}_{\mathcal{A}} \models \text{PSIZE}$*

Fact 4.1.5. *$\text{PSIZE} \not\models \text{PCS}$.*

Proof. PSIZE fails to axiomatize PCS for the same reason that BASIC fails to axiomatize CS: the lack of divisibility principles. $\square$

We offer PCA as an axiomatic version of PCS:

Definition 4.1.6. *PCA, the pure class size axioms, is defined as:*

$$\text{PSIZE} \cup \{ \text{ADIV}_n \mid n > 0 \}$$

Fact 4.1.7. *If $\mathcal{A} \models \text{CA}$, then $\mathcal{S}_{\mathcal{A}} \models \text{PCA}$*

If $\mathcal{A}$ is a standard finite interpretation of BASIC with n atoms, then the elements of $\mathcal{S}_{\mathcal{A}}$ can be regarded as the sequence $0, \dots, n$ with the usual ordering, where

$$\mathcal{S}_{\mathcal{A}} \models \text{Unit}(x) \text{ iff } x = 1$$

and

$$\mathcal{S}_{\mathcal{A}} \models \text{Sum}(i, j, k) \text{ iff } (i + j) = k \leq n$$

Once n is fixed, this is the only interpretation allowed by the axioms PSIZE. In particular, MONOT rules out the interpretation in which $\text{Sum}(i, j, k)$ is satisfied just in case $(i + j) \equiv k \pmod{n + 1}$.

4.2 Some models and theories

We shall use Theorem 4.2.1a to show that PCA is model complete and Theorem 4.2.1b to show, in chapter 5, that CAI is model complete.

Theorem 4.2.1.

- a. If T satisfies **Monk's Condition**, then T is model complete.

Monk's Condition. If $\mathcal{A} \models T$, $\mathcal{B} \models T$, $\mathcal{A} \subseteq \mathcal{B}$, and $\mathcal{C}$ is a finitely generated submodel of $\mathcal{B}$, then there is an isomorphism, $f: \mathcal{C} \rightarrow \mathcal{A}$ such that if $x \in \mathcal{C} \cap \mathcal{A}$, then $f(x) = x$. f is a **Monk mapping**.

- b. If T is model complete and L_T has no function symbols, then T satisfies Monk's Condition.

Proof.

- a. [Monk, p.359]
- b. If L_T has no function symbols, then any finitely generated structure over L_T is finite. So, suppose $\mathcal{C}$ contains $a_1, \dots, a_n$ (from $\mathcal{A}$) and $b_1, \dots, b_m$ (from $\mathcal{B} - \mathcal{A}$). Let ϕ_1 be the diagram of $\mathcal{C}$ and obtain ϕ_2 from ϕ_1 by substituting the variable x_i for each constant a_i and the variable y_i for each constant b_i . Finally, let ϕ_3 be

$$\exists y_1 \dots \exists y_m \phi_2$$

So, ϕ_3 is a primitive formula. $\mathcal{B} \models \phi_3(a_1, \dots, a_n)$, so $\mathcal{A}$ does as well, by Fact C.5d in the Appendix. So, to obtain the desired isomorphism, map the a_i 's into themselves and map the b_i 's into a sequence of elements of $\mathcal{A}$ which can stand in for the existentially quantified variables of ϕ_3 .

□

In chapter 5, we use Monk's Theorem to infer the model completeness of the theory CA from that of PCA. To establish the model completeness of PCA, we use Fact 4.2.2.

Fact 4.2.2.

- a. If T_1 is model complete and $T_1 \vdash T_2$, then T_2 is also model complete.
- b. If T is model complete in L , and L' is an expansion of L by adjoining new individual constants, then T is model complete in L' . [Monk, p.355]

Definition 4.2.3. Suppose L_1 and L_2 are first order languages and $L_{12} = L_1 - L_2$. A **translation (simple translation)** of L_1 into L_2 is a function, τ , which:

- a. assigns to the universal quantifier a (quantifier free) formula, $\tau_{\forall}$, of L_2 with exactly one free variable.
- b. assigns to each n -place predicate, P , in L_{12} a (quantifier free) formula, τ_P , of L_2 with exactly n free variables, and
- c. assigns to each n -place function symbol, O , in L_{12} a (quantifier free) formula, τ_O , of L_2 with exactly $(n + 1)$ free variables.

Definition 4.2.4. If τ is a translation of L_1 into L_2 , then τ extends to all formulae of L_1 as follows:

- a. Predicates and function symbols of L_2 are translated into themselves.
- b.

$$\begin{aligned}
 \tau(\phi \vee \psi) &= \tau(\phi) \vee \tau(\psi) \\
 \tau(\phi \wedge \psi) &= \tau(\phi) \wedge \tau(\psi) \\
 \tau(\neg\phi) &= \neg\tau(\phi) \\
 \tau(\forall x\phi) &= \forall x(\tau_{\forall}(x) \rightarrow \phi) \\
 \tau(\exists x\phi) &= \exists x(\tau_{\forall}(x) \wedge \phi)
 \end{aligned}$$

Definition 4.2.5. If τ is a translation of L_1 into L_2 , then

- a. The **functional assumptions** of τ are the sentences:

$$\begin{aligned}
 \forall x_1 \dots \forall x_n (\tau_{\forall}(x_1) \wedge \dots \wedge \tau_{\forall}(x_n)) \rightarrow \\
 \exists y_1 (\tau_{\forall}(y_1) \wedge \forall y_2 (\tau_O(x_1, \dots, x_n, y_2) \leftrightarrow y_1 = y_2))
 \end{aligned}$$

where O is a function symbol in L_1 but not in L_2 .

- b. The **existential assumption** of τ is

$$\exists x \tau_{\forall}(x)$$

The functional assumptions of a translation say that the formulas which translate function symbols yield unique values within the relevant part of the domain when given values in the relevant part of the domain. The *relevant part of the domain* is the set of elements which satisfy the interpretation of universal quantifier. The existential assumption of a translation says that that subdomain is non-empty. Notice that the existential and functional assumptions of a translation are sentences of L_2 .

A translation from L_2 into L_1 induces a mapping from interpretations of L_2 into interpretations of L_1 .

Definition 4.2.6. If τ is a translation from L_1 into L_2 and $\mathcal{B}$ is an interpretation of L_2 which satisfies the existential and functional assumptions of τ , then $\tau(\mathcal{B})$ is the interpretation $\mathcal{A}$ of L_1 such that:

- a. $\mathcal{A}$ is the set of elements of $\mathcal{B}$ which satisfy $\tau_{\forall}$.
- b. $\mathcal{A}$ interprets all predicates and function symbols common to L_1 and L_2 in the same way that $\mathcal{B}$ does.
- c. $\mathcal{A}$ interprets all predicates and function symbols in L_{12} in accordance with the translations assigned by τ .

$$\begin{aligned}\mathcal{A} \models P(\vec{x}) & \text{ iff } \mathcal{B} \models \tau_P(\vec{x}) \\ \mathcal{A} \models y = O(\vec{x}) & \text{ iff } \mathcal{B} \models \tau_O(\vec{x}, y)\end{aligned}$$

The following condition on theories allows us to infer the model completeness of one from the model completeness of the other.

Definition 4.2.7.

- a. If τ is a translation from L_1 to L_2 , then T_1 is **τ -reducible** to T_2 iff for every model $\mathcal{A}$ of T_1 there is a model $\mathcal{B}$ of T_2 such that $\mathcal{A} = \tau(\mathcal{B})$.
- b. T_1 is **reducible (simply reducible)** to T_2 iff there is a (simple) translation, τ , for which T_1 is τ -reducible to T_2 .
- c. T_1 is **uniformly τ -reducible** to T_2 iff for any models, $\mathcal{A}_1$ and $\mathcal{B}_1$, such that $\mathcal{A}_1 \models T_1$ and $\mathcal{B}_1 \models T_1$, and $\mathcal{A}_1 \subseteq \mathcal{B}_1$, there exist models $\mathcal{A}_2$ and $\mathcal{B}_2$ such that $\mathcal{A}_2 \models T_2$ and $\mathcal{B}_2 \models T_2$, and $\mathcal{A}_2 \subseteq \mathcal{B}_2$ $\mathcal{A}_1 = \tau(\mathcal{A}_2)$ and $\mathcal{B}_1 = \tau(\mathcal{B}_2)$,

Lemma 4.2.8. Suppose that T_1 is τ -reducible to T_2 and that $\mathcal{A}_1 = \tau(\mathcal{A}_2)$. Then, for any primitive formula, ϕ , of L_1 and any sequence, $\vec{x} \in \mathcal{A}_1$,

$$\mathcal{A}_1 \models \phi(\vec{x}) \text{ iff } \mathcal{A}_2 \models \tau_{\phi}(\vec{x})$$

Proof. Suppose

$$\phi(\vec{x}) = \exists y_1 \dots \exists y_n \phi'(\vec{x})$$

where $\phi'(\vec{x})$ is a conjunction of atomic formulae and negations of atomic formulae. Then

$$\begin{aligned}\mathcal{A}_1 \models \phi(\vec{x}) & \text{ iff } \mathcal{A}_1 \models \exists y_1 \dots \exists y_n \phi'(\vec{x}, y_1, \dots, y_n) \\ & \text{ iff } \mathcal{A}_1 \models \phi'(\vec{x}, b_1, \dots, b_n), \text{ for } b_i \in \mathcal{A} \\ & \text{ iff } \mathcal{A}_2 \models \tau_{\phi'}(\vec{x}, b_1, \dots, b_n) \\ & \text{ iff } \mathcal{A}_2 \models \exists y_1 \dots \exists y_n \\ & \quad (\tau_{\forall}(y_1) \wedge \dots \wedge \tau_{\forall}(y_n) \wedge \phi'(\vec{x}, y_1, \dots, y_n)) \\ & \text{ iff } \mathcal{A}_2 \models \tau_{\phi}(\vec{x})\end{aligned}$$

□

Theorem 4.2.9. *If T_2 is model complete, τ is a simple translation from L_{T_1} to L_{T_2} , and T_1 is uniformly τ -reducible to T_2 , then T_1 is also model complete.*

Proof. By Fact C.5d in the Appendix, it is enough to show that given models $\mathcal{A}_1$ and $\mathcal{B}_1$ of T_1 , where $\mathcal{A}_1 \subseteq \mathcal{B}_1$ and a primitive formula, ϕ :

$$\begin{array}{ll} \text{If} & \mathcal{B}_1 \models \phi(\vec{x}) \text{ for } \vec{x} \in \mathcal{A} \\ \text{then} & \mathcal{A}_1 \models \phi(\vec{x}) \end{array}$$

Since T_1 is uniformly τ -reducible to T_2 , there are models of T_2 , $\mathcal{A}_2 \subseteq \mathcal{B}_2$, where $\mathcal{A}_1 = \tau(\mathcal{A}_2)$ and $\mathcal{B}_1 = \tau(\mathcal{B}_2)$

$$\begin{array}{ll} \text{Since} & \mathcal{B}_1 \models \phi(\vec{x}) & \text{by assumption} \\ \text{then} & \mathcal{B}_2 \models \tau_\phi(\vec{x}) & \text{by lemma 4.2.8} \\ \text{So} & \mathcal{A}_2 \models \tau_\phi(\vec{x}) & \text{since } T_2 \text{ is model complete} \\ \text{and} & \mathcal{A}_1 \models \phi(\vec{x}) & \text{by lemma 4.2.8} \end{array}$$

□

We shall now define several theories, all more or less familiar, which will serve in showing that our theory of size is model complete.

Definition 4.2.10.

a. The theory of abelian groups with identity has the following axioms

$$\begin{array}{ll} (1) & x + (y + z) = (x + y) + z \\ (2) & x + y = y + x \\ (3) & x + 0 = x \\ (4) & \exists y(x + y = 0) \end{array}$$

b. The theory of cancellable abelian semigroups with identity consists of (1), (2), and (3) above and:

$$(4') \quad x + y = x + z \rightarrow y = z$$

c. The axioms of simple order are:

$$\begin{array}{ll} (5) & x \leq y \wedge y \leq z \rightarrow x \leq z \\ (6) & x \leq y \wedge y \leq x \rightarrow x = y \\ (7) & x \leq x \\ (8) & x \leq y \vee y \leq x \end{array}$$

d. The theory of $\mathbb{Z}$ -groups, $\mathbf{Zg}$, has the following axioms

(i) The axioms for abelian groups with identity,

- (ii) The axioms for simple order,
- (iii) The following additional axioms:

$$(9) \quad y \leq z \rightarrow x + y \leq x + z$$

$$(10) \quad 1 \text{ is the least element greater than } 0$$

$$(11) \quad \forall x \exists y (ny = x \vee \dots \vee ny = x + (n - 1))$$

for each positive n , where ny stands for:

$$\underbrace{y + \dots + y}_{n \text{ times}}$$

e. The **theory of $\mathbb{N}$ -semigroups** has the following axioms:

- (i) The axioms for cancellable abelian semigroups with identity,
- (ii) The axioms for simple order,
- (iii) Axioms (9) and (10) of Zg , and
- (iv) The additional axiom:

$$(12) \quad 0 \leq x$$

f. The **theory of $\mathbb{Z}$ -groups modulo I** , Zgm , consists of the following axioms:

- (i) The axioms for abelian groups with identity,
- (ii) The axioms for simple order
- (iii) Axioms (10) and (11) of Zg , axiom (12) from the theory of $\mathbb{N}$ -semigroups, and
- (iv) The following additional axioms

$$(13) \quad (y \leq z \wedge x \leq x + z) \rightarrow x + y \leq x + z$$

$$(14) \quad x \leq I$$

The theory of $\mathbb{Z}$ -groups is taken from [Chang, p. 291]

Fact 4.2.11.

- a. Zg is the complete theory of $\langle \mathbb{Z}, +, 0, 1, \leq \rangle$. [Chang, p.291]
- b. Zg is model complete.[Robinson]

Theorem 4.2.12.

- a. The theory of $\mathbb{N}$ -semigroups is model complete.

b. *The theory of $\mathbb{Z}$ -groups modulo I is model complete.*

Proof.

- a. Every abelian semigroup with cancellation can be isomorphically embedded in an abelian group [Kurosh, pp. 44-48]. It is clear from the construction in [Kurosh] that if the semigroup is ordered, the abelian group in which it is embedded may also be ordered and that the elements of the semigroup will be the positive elements of the group. Moreover, the (rough) divisibility of the elements in the semigroup will be carried over to the group.

Consequently, the theory of $\mathbb{N}$ -semigroups is uniformly reducible to the theory of $\mathbb{Z}$ groups by the translation:

$$\tau_V = \ulcorner 0 \leq x \urcorner$$

Since the latter is model complete, so is the former, by 4.2.9.

- b. First, consider the theory of $\mathbb{N}$ -semigroups in the language which contains, besides the constant symbols in the original theory, an individual constant, I . The theory of $\mathbb{N}$ -semigroups is model-complete in this language, by 4.2.2b.

We claim that the theory of $\mathbb{Z}$ -groups modulo I is uniformly reducible to this new theory by the following translation:

$$\begin{aligned} \tau_V &= \ulcorner x \leq I \urcorner \\ \tau_+ &= \ulcorner x + y = z \vee x + y = I + z \urcorner \end{aligned}$$

(The construction: given a model of Zgm , stack up ω many copies of the model, assigning interpretations in the obvious way. The result is an $\mathbb{N}$ -semigroup and the original model is isomorphic to the first copy of itself.)

□

In the next section, we use Theorem 4.2.9 to show that PSIZE is model complete, by reducing it to the theory of $\mathbb{Z}$ -groups modulo I . Chapter 5 uses Monk's Theorem to show that CA is model complete.

4.3 PCA is model complete

To show that PCA is model complete, we shall reduce it to Zgm , the theory of $\mathbb{Z}$ -groups with addition taken modulo some constant (see 4.2.11). The model completeness of PCA then follows from the model completeness of Zgm (Fact 4.2.12) and Theorem 4.2.9. Specifically, we shall show that every model of PCA is the τ -image of a model of Zgm , where τ is the following translation.

Definition 4.3.1. Let τ be the translation from L_{PSIZE} to L_{Zgm} where:

$$\begin{aligned}\tau_{\forall} &= \ulcorner x = x \urcorner \\ \tau_{Sum(x,y,z)} &= \ulcorner x + y = z \wedge x \leq z \urcorner \\ \tau_{Unit(x)} &= \ulcorner x = 1 \urcorner \\ \tau_{\tilde{x}} &= \ulcorner x + y = \mathbf{I} \urcorner \\ \tau_{x < y} &= \ulcorner x \leq y \wedge x \neq y \urcorner \\ \tau_{\emptyset} &= \ulcorner x = 0 \urcorner\end{aligned}$$

Given a model, $\mathcal{A}$, of PCA, we can construct a model, $\mathcal{Z}_{\mathcal{A}}$, of Zgm directly:

Definition 4.3.2. If $\mathcal{A} \models PSIZE$, then $\mathcal{Z}_{\mathcal{A}}$ is the interpretation of L_{Zgm} in which:

- (a) $\dot{\mathcal{Z}}_{\mathcal{A}} = \dot{\mathcal{A}}$
- (b) $\mathcal{Z}_{\mathcal{A}} \models (x \leq y)$ iff $\mathcal{A} \models (x < y \vee x = y)$
- (c) $\mathcal{Z}_{\mathcal{A}}(I) = \mathcal{A}(I)$
- (d) $\mathcal{Z}_{\mathcal{A}}(0) = \mathcal{A}(0)$
- (e) $\mathcal{Z}_{\mathcal{A}} \models (x + y = z)$ iff $\mathcal{A} \models Sum(x, y, z)$
or $\mathcal{A} \models \exists a \exists w Unit(a)$
 $\wedge Sum(\tilde{x}, \tilde{y}, w) \wedge Sum(w, a, \tilde{z})$
- (f) $\mathcal{Z}_{\mathcal{A}} \models (x = 1)$ iff $\mathcal{A} \models Unit(x)$

Fact 4.3.3j establishes that 4.3.2 gives Sum a functional interpretation. Theorem 4.3.8 establishes that $\mathcal{Z}_{\mathcal{A}} \models Zgm$, on the basis of the intervening facts: 4.3.3 deals with the model $\mathcal{A}$ of PSIZE; 4.3.4 deals with the corresponding model $\mathcal{Z}_{\mathcal{A}}$; 4.3.6 verifies some connections between $\mathcal{A}$ and $\mathcal{Z}_{\mathcal{A}}$.

Fact 4.3.3. The following are theorems of PSIZE. ($Sm(x, y)$ abbreviates $\exists z Sum(x, y, z)$.)

- (a) $Sum(x, y, z_1) \wedge Sum(x, y, z_2) \rightarrow z_1 = z_2$ ($UNIQ^+$)
- (b) $Sum(x, y_1, z) \wedge Sum(x, y_2, z) \rightarrow y_1 = y_2$ ($UNIQ^-$)
- (c) $Sum(x, y, w_1) \ \& \ Sum(w_1, z, w) \rightarrow$ ($ASSOC2$)
 $\exists w_2 (Sum(y, z, w_2) \wedge Sum(x, w_2, w))$
- (d) $\tilde{x} = x$
- (e) $x < y \leftrightarrow \tilde{y} < \tilde{x}$
- (f) $x < y \wedge y < \tilde{y} \rightarrow x < \tilde{x}$
- (g) $Sm(x, y) \vee Sm(\tilde{x}, \tilde{y})$
- (h1) $Sum(x, y, z) \rightarrow Sum(\tilde{z}, y, \tilde{x})$
- (h2) $Sum(x, y, z) \rightarrow Sum(x, \tilde{z}, \tilde{y})$
- (i) $Sum(x, y, z_1) \wedge Sum(\tilde{x}, \tilde{y}, \tilde{z}_2) \rightarrow (z_1 = z_2 = I)$
- (j) $\exists z_1 Sum(x, y, z_1) \leftrightarrow$
 $\neg \exists a \exists w \exists z_2 (Unit(a) \wedge Sum(\tilde{x}, \tilde{y}, w) \wedge Sum(w, a, \tilde{z}_2))$

Proof. The proofs are elementary. $\square$

In addition to the theorems of PSIZE listed in 4.3.3, we require a battery of tedious facts about the model $\mathcal{Z}_{\mathcal{A}}$. We shall state these in terms of the model $\widehat{\mathcal{Z}}_{\mathcal{A}}$, an expansion of both $\mathcal{A}$ and $\mathcal{Z}_{\mathcal{A}}$, which interprets two additional operators, as follows:

Definition 4.3.4. *Given a model $\mathcal{A}$ of PSIZE, $\widehat{\mathcal{Z}}_{\mathcal{A}}$ is the expansion of $\mathcal{Z}_{\mathcal{A}}$ induced by the definitions in 4.3.2 together with (a) and (b):*

- (a) $\widehat{\mathcal{Z}}_{\mathcal{A}} \models y = -x$ iff $\mathcal{Z}_{\mathcal{A}} \models x + y = 0$
- (b) $\widehat{\mathcal{Z}}_{\mathcal{A}} \models z = x - y$ iff $\mathcal{Z}_{\mathcal{A}} \models z = x + -y$

Remark. *Given a model $\mathcal{A}$ of PSIZE: for each x in $\dot{\mathcal{Z}}_{\mathcal{A}}$, there is a unique y such that:*

$$\mathcal{Z}_{\mathcal{A}} \models x + y = 0$$

Proof.

- a. If $x = 0$, then $\mathcal{Z}_{\mathcal{A}} \models x + y = 0$ iff $y = 0$.

$\Rightarrow$

If	$\mathcal{Z}_{\mathcal{A}} \models 0 + y = 0$	
then	$\mathcal{A} \models Sum(\emptyset, y, \emptyset)$	by 4.3.2e
But	$\mathcal{A} \models Sum(\emptyset, \emptyset, \emptyset)$	by IDENT
So	$\mathcal{Z}_{\mathcal{A}} \models y = 0$	by $UNIQ^-$

$\Leftarrow$ Obvious

b. If $x > 0$, there is a unique y which satisfies

$$(*) \quad \text{Sum}(\tilde{x}, a, y), \text{ where Unit } (a)$$

since there is a unique unit and $\mathcal{A} \models \text{UNIQ}^+$.

But $\mathcal{Z}_{\mathcal{A}} \models x + y = 0$ iff $\mathcal{A} \models (*)$.

$\Rightarrow$

If	$\mathcal{Z}_{\mathcal{A}} \models x + y = 0$	
then	$\mathcal{A} \models \text{Sum}(\tilde{x}, \tilde{y}, w)$	
and	$\mathcal{A} \models \text{Sum}(w, a, \tilde{\emptyset})$	since $\mathcal{A} \not\models \text{Sum}(x, y, 0)$
But	$\mathcal{A} \models \tilde{\emptyset} = \mathbf{I}$	
So	$\mathcal{A} \models w = \tilde{a}$	by COMP and UNIQ^-
So	$\mathcal{A} \models \text{Sum}(\tilde{x}, \tilde{y}, \tilde{a})$	
So	$\mathcal{A} \models \text{Sum}(\tilde{x}, a, y)$	by 4.3.3d and h2

$\Leftarrow$

If	$\mathcal{A} \models (*)$	
then	$\mathcal{A} \models \text{Sum}(\tilde{x}, \tilde{y}, \tilde{a})$	by 4.3.3h2
and	$\mathcal{A} \models \text{Sum}(\tilde{a}, a, \tilde{\emptyset})$	by COMP
So	$\mathcal{Z}_{\mathcal{A}} \models x + y = 0$	by 4.3.2e

□

Fact 4.3.5. $\widehat{\mathcal{Z}}_{\mathcal{A}}$ satisfies the following:

- | | |
|-----|---|
| (a) | $-(x + y) = -x + -y$ |
| (b) | $-(x - y) = y - x$ |
| (c) | $(x + y) - y = x$ |
| (d) | $-(-x) = x$ |
| (e) | $x \neq 0 \wedge x < y \rightarrow -y < -x$ |

Proof. Omitted.

□

Fact 4.3.6. $\widehat{\mathcal{Z}}_{\mathcal{A}}$ satisfies the following:

- a. $\text{Sm}(a, b) \leftrightarrow \text{Sum}(a, b, a + b)$
- b. $c \leq b \leftrightarrow \text{Sum}(c, b - c, b)$

- c. $Sum(a, b, c) \wedge 0 < b \rightarrow Sum(a, -c, -b)$
 $Sum(a, b, c) \wedge 0 < a \rightarrow Sum(-c, b, -a)$
- d. $\neg Sm(a, b) \rightarrow (Sm(-a, -b) \vee b = -a)$

Proof.

a. $(\Rightarrow)$

Suppose $Sm(a, b)$
 So $Sum(a, b, w)$ for some w
 So $(a + b) = w$ by 4.3.2e
 So $Sum(a, b, a + b)$

$(\Leftarrow)$ Obvious.

b. $(\Rightarrow)$

Suppose $c \leq b$
 then $Sum(c, w, b)$ for some w by EXIST⁻
 So $b = c + w$ by (a) and UNIQ⁺
 So $b - c = (c + w) - c$
 So $b - c = w$ by 4.3.5c
 So $Sum(c, b - c, b)$

$(\Leftarrow)$

Suppose $Sum(c, b - c, b)$
 But $Sum(c, 0, c)$ by IDENT
 So $0 \leq (b - c) \leftrightarrow b \leq c$ by MONOT and UNIQ⁺
 So $b \leq c$ by MIN

c.

Suppose $Sum(a, b, c)$
 and $0 < b$
 then $c = (a + b)$ by (a) and UNIQ⁺
 So $c - b = (a + b) - b$
 So $c - b = a$ by 4.3.5c
 So $-b = a - c$ by 4.3.5c
 So $Sum(a, -c, -b)$ if $Sm(a, -c)$ by (a)
 But if $a = 0$
 then $Sm(a, -c)$ by IDENT

And if $a \neq 0$
 then $a < c$, since $0 < b$ by MONOT and UNIQ⁺
 So $-c < -a$ by 4.3.5e
 So $-c \leq \tilde{a}$ by 4.3.4
 But $\text{Sm}(a, \tilde{a})$ by IDENT
 So $\text{Sm}(a, -c)$ by EXIST⁺

d.

Suppose $\neg \text{Sm}(a, b)$
 But $b = (a + b) - a$, by 4.3.5c
 So $\neg \text{Sum}(a, (a + b) - a, a + b)$, by (a)
 So $\neg(a \leq a + b)$, by (b)
 So $(a + b) < a$, by TRICH
 But if $(a + b) \neq 0$
 then $-a < -(a + b)$, by 4.3.5e
 So $-a < -a + -b$, by 4.3.5a
 So $\text{Sum}(-a, (-a + -b) - (-a), -a + -b)$, by (b)
 But $(-a + -b) - (-a) = -b$, by 4.3.5c
 So $\text{Sum}(-a, -b, -a + -b)$
 So $\text{Sm}(-a, -b)$
 And if $(a + b) = 0$
 then $b = -a$

□

We can now show that $\mathcal{Z}_A \models \text{Zgm}$. The only real difficulty arises in verifying that addition is associative. We need the following lemma.

Lemma 4.3.7. $\mathcal{Z}_A \models (a + c) + (b - c) = a + b$

Proof. We shall work through successively more general cases:

a. When $\text{Sm}(a, b)$ and $c \leq b$

We know $\text{Sum}(b - c, c, b)$ by 4.3.6b
 and $\text{Sum}(b, a, a + b)$ by 4.3.6a
 So $\exists w(\text{Sum}(c, a, w)$
 $\quad \wedge \text{Sum}(b - c, w, a + b))$ by ASSOC2
 So $w = c + a$ by 4.3.6a and UNIQ⁺
 and $a + b = (b - c) + (c + a)$ by 4.3.6a and UNIQ⁺
 So $a + b = (a + c) + (b - c)$

b. When $\text{Sm}(a, b)$ and $\text{Sm}(a, c)$

If	$c \leq b$, Case (a) applies directly	
So assume	$b < c$	
then	$a + c = (a + b) + (c - b)$	by Case (a)
So	$(a + c) - (c - b) = a + b$	by 4.3.5c
So	$(a + c) + -(c - b) = a + b$	by 4.3.4b
So	$(a + b) + (b - c) = a + b$	by 4.3.5b

c. $\text{Sm}(a, b)$

If	$\text{Sm}(a, c)$ Case (b) applies	
So assume	$\neg \text{Sm}(a, c)$ and thus $b < c$	by EXIST^+
So	$\text{Sum}(b, c - b, c)$	by 4.3.6b
and	$0 < c - b$	by MONOT
So	$\text{Sum}(b, -c, -(c - b))$	by 4.3.6c
So	$\text{Sum}(b, -c, b - c)$	by 4.3.5b
So	$\text{Sum}(-c, b, b - c)$	(c1)
Either	$\text{Sm}(-a, -c)$ or $c = -a$ since $\neg \text{Sm}(a, c)$	by 4.3.6d
Assume	$\text{Sm}(-a, -c)$	
then	$\text{Sum}(-a, -c, -a + -c)$	by 4.3.6a
So	$\text{Sum}(-a, -c, -(a + c))$	by 4.3.5a
and	$0 < -a$	
So	$\text{Sum}(-(-(a + c)), -c, -(-a))$	by 4.3.6c
So	$\text{Sum}(a + c, -c, a)$	by 4.3.5d (c2)
Assume	$c = -a$	
then	$a + c = 0$	
and	$a = -c$	
Hence	$\text{Sum}(a + c, -c, a)$	by (c2)
and	$\text{Sum}(-c, b, b - c)$	by (c1)
and	$\text{Sum}(a, b, a + b)$	since $\text{Sm}(a, b)$
So	$\text{Sum}(a + c, b - c, a + b)$	by ASSOC
So	$(a + c) + (b - c) = a + b$	by 4.3.6a

d. Whenever

Suppose $\neg \text{Sm}(a, b)$, for otherwise case c applies.

Either	$\text{Sm}(-a, -b)$ or $b = -a$	by 4.3.6d
Assume	$b = -a$	
then	$a + b = 0$	
and	$b - c = (-a) - c$	
	$= -a + -c$	
	$= -(a + c)$	
So	$(a + b) + (b - c)$	
	$= (a + c) + -(a + c)$	
	$= 0 = a + b$	
Assume	$\text{Sm}(-a, -b)$	
then	$-a + -b = (-a + -c) + (-b - -c)$	by case c
So	$-(a + b) = -(a + c) + -(b - c)$	by 4.3.5a
So	$-(a + b) = -((a + c) + (b - c))$	by 4.3.5a
So	$a + b = (a + c) + (b - c)$	by 4.3.5d

□

Theorem 4.3.8. $\mathcal{Z}_{\mathcal{A}} \models \text{Zgm}$

Proof. The only axiom for abelian groups that needs further verification is associativity. We prove this using lemma 4.3.7:

$$\begin{aligned} x + (y + z) &= (x + y) + ((y + z) - y), \text{ by 4.3.7} \\ &= (x + y) + z, \text{ by 4.3.5c} \end{aligned}$$

The ordering axioms of Zgm are satisfied in $\mathcal{Z}_{\mathcal{A}}$ because $\mathcal{Z}_{\mathcal{A}}$ uses the same ordering as $\mathcal{A}$, $\mathcal{A} \models \text{PSIZE}$, and PSIZE includes the same ordering axioms.

$\mathcal{Z}_{\mathcal{A}}$ satisfies axiom (10) of Zgm because $\mathcal{A}$ satisfies the UNIT axiom of PSIZE.

The divisibility of elements in $\mathcal{Z}_{\mathcal{A}}$ required by axiom (11) of Zgm is guaranteed by the fact that $\mathcal{A}$ satisfies the divisibility principles of PCA.

Similarly, $\mathcal{Z}_{\mathcal{A}}$ satisfies axioms (12), and (14) of Zgm because $\mathcal{A}$ satisfies MIN and MAX.

Axiom (13) of Zgm is:

$$y \leq z \wedge x \leq x + z \rightarrow x + y \leq x + z$$

If	$\mathcal{Z}_{\mathcal{A}} \models x \leq x + z$	
then	$\mathcal{A} \models \text{Sum}(x, z, x + z)$	
Since	$\mathcal{Z}_{\mathcal{A}} \models y < z$	
then	$\mathcal{A} \models \text{Sum}(x, y, x + y)$	by EXIST ⁺
and	$\mathcal{A} \models x + y \leq x + z$	by MONOT
So	$\mathcal{Z}_{\mathcal{A}} \models x + y \leq x + z$	

□

So, we PCA is τ -reducible to Zgm. The reduction is uniform since each model $\mathcal{A}$ of PCA has the same domain as its Zgm-model. So, we may conclude that PCA is model complete.

Theorem 4.3.9.

- a. *PCA is model complete.*
- b. *PCA satisfies Monk's condition.*

Proof.

- a. Apply Theorem 4.2.9.
- b. Immediate from (a) and 4.2.1(b).

□

4.4 Remarks on showing that PCA axiomatizes PCS

To prove that $\text{PCA} \equiv \text{PCS}$, we could follow the method outlined at the end of chapter 3 for showing that $\text{CA} \equiv \text{CS}$.

We already know that $\text{PCS} \vdash \text{PCA}$, so we need only prove (1), which follows from (2) by 3.3.1b.

- (1) $\text{PCA} \vdash \text{PCS}$
- (2) Every completion of PCA is consistent with PCS.

But (2) is equivalent to the conjunction of (2a) and (2b).

- (2a) Every finite completion of PCA is consistent with PCS.
- (2b) Every infinite completion of PCA is consistent with PCS.

The finite completions of PCA are the theories $\text{PCA}; \text{EXACTLY}_n$. But $\text{PCA}; \text{EXACTLY}_n$ is true in $\mathcal{S}_{\mathcal{A}}$, where $\mathcal{A}$ is the standard finite interpretation of $L_{C<}$ containing n atoms. So, the finite completions of PCA are consistent with PCS. (Formally, we would have to define EXACTLY_n in terms of units rather than atoms.)

Letting $\text{PCAI} = \text{PCA} + \text{INF}$ and $\text{PCSI} = \text{PCS} + \text{INF}$, (2b) is a consequence of (3a) and (3b).

- (3a) If PCAI_f is consistent, then PCSI_f is consistent.
- (3b) If T is a completion of PCAI, then $T \equiv \text{PCAI}_f$, for some f .

To prove (3a), we would have to prove analogues of 3.3.9 through 3.3.13 for PCA and PCS. This seems straightforward, but tedious. The trick is to show that enough axioms about size have been incorporated in PCA to establish the entailments among MOD statements.

To prove (3b), it is sufficient to demonstrate (4), because every completion of PCAI entails PCAI_f for some total f .

(4) If f is total, then PCAI_f is complete.

But PCA is model complete, so only (5) remains to be shown.

(5) If f is total and congruous, then PCAI_f has a prime model.

We will not construct prime models for the extensions of PCAI. It's apparent that the size models of the prime models for CAI_f would do nicely. Alternatively, the construction could be duplicated in this simpler case.

5

Completeness of CA

5.1 Model completeness of CA

We shall show that CA is model complete by showing that it satisfies Monk's criterion (see 4.3.1). So, given Assumption 5.1.1, we want to prove 5.1.2.

Assumption 5.1.1. $\mathcal{A} \models CA$, $\mathcal{B} \models CA$, $\mathcal{A} \subseteq \mathcal{B}$, and $\mathcal{C}$ is a finitely generated substructure of $\mathcal{B}$.

Theorem 5.1.2. *There is an isomorphic embedding, $f: \mathcal{C} \rightarrow \mathcal{A}$, where*

$$f(x) = x \text{ if } x \in \dot{\mathcal{C}} \cap \dot{\mathcal{A}}$$

The Monk mappings for PCA can serve as a guide in constructing Monk mappings for CA. The existence of Monk mappings for PCA tells us that we can find elements with the right sizes. $\text{DISJ}^\cup$ and $\text{REP}^<$ then allow us to find elements with those sizes that fit together in the right way.

Strictly speaking, $\mathcal{S}_{\mathcal{A}}$ is not a submodel of $\mathcal{S}_{\mathcal{B}}$, so we cannot apply Monk's Theorem directly. But, let

$$\mathcal{S}_{\mathcal{B}}(X) = \text{the submodel of } \mathcal{S}_{\mathcal{B}} \text{ whose domain is } \{ \sigma_{\mathcal{B}}[x] \mid x \in X \}$$

Then, clearly,

$$\mathcal{S}_{\mathcal{A}} \simeq \mathcal{S}_{\mathcal{B}}(A) \subseteq \mathcal{S}_{\mathcal{B}}, \text{ and}$$

$$\mathcal{S}_{\mathcal{C}} \simeq \mathcal{S}_{\mathcal{B}}(C), \text{ a finitely generated submodel of } \mathcal{S}_{\mathcal{B}}.$$

Monk's Theorem applies directly to $\mathcal{S}_{\mathcal{B}}(A)$, $\mathcal{S}_{\mathcal{B}}$, and $\mathcal{S}_{\mathcal{B}}(C)$, so we may conclude 5.1.3:

Fact 5.1.3. *There is an isomorphic embedding, $g: \mathcal{S}_{\mathcal{C}} \rightarrow \mathcal{S}_{\mathcal{A}}$, where*

$$g(\sigma_{\mathcal{B}}[x]) = \sigma_{\mathcal{A}}[x] \text{ for all } x \in \dot{\mathcal{C}} \cap \dot{\mathcal{A}}.$$

That is to say, the sizes of elements in $\mathcal{C}$ can be embedded in the sizes of elements in $\mathcal{A}$. It remains to be shown that the elements of $\mathcal{C}$ themselves can be mapped into $\mathcal{A}$ by a function which preserves boolean relations as well as size relations.

$\mathcal{C}$ is a finite, and hence atomic, boolean algebra, though its atoms need not be atoms of $\mathcal{B}$; indeed, if $\mathcal{B}$ is infinite, there must be some atoms of $\mathcal{C}$ which are not atoms of $\mathcal{B}$ since the union of all atoms of $\mathcal{C}$ is the basis of $\mathcal{B}$.

Definition 5.1.4. d is a **molecule** iff $d \in \dot{\mathcal{C}} \cap \dot{\mathcal{A}}$ and no proper subset of d is in $\dot{\mathcal{C}} \cap \dot{\mathcal{A}}$.

$\dot{\mathcal{C}} \cap \dot{\mathcal{A}}$ is a boolean algebra whose basis is the same as the basis of $\mathcal{C}$. So every atom of $\mathcal{C}$ is included in some molecule. The embedding f has to map each molecule to itself. Moreover, f has to be determined by its values on $\mathcal{C}$ since f must preserve unions. In fact, the atoms of $\mathcal{C}$ can be partitioned among the molecules. So, if

$$d = b_1 \cup \dots \cup b_n$$

where d is a molecule and $b_1, \dots, b_n$ are the atoms of $\mathcal{C}$ contained in d , then d must also be the (disjoint) union of $f(b_1), \dots, f(b_n)$. If this condition is satisfied, f will preserve boolean relations. f must also select images with appropriate sizes.

Proof. Proof of 5.1.2 Given a molecule, d , let $b_1, \dots, b_n$ be the atoms of $\mathcal{C}$ contained in d . For each b_i , let c_i be some member of $g(\sigma_{\mathcal{B}}[b_i])$. These elements will be elements of $\dot{\mathcal{A}}$, since g yields sizes in $\mathcal{A}$ whose members are in $\dot{\mathcal{A}}$. These elements have the right sizes, as we will show below, but they are in the wrong places. We have no guarantee that they are contained in the molecule d . So we still need to show that there are disjoint elements of $\dot{\mathcal{A}}$, $a_1, \dots, a_n$ whose union is d and whose sizes are the same as those of $b_1, \dots, b_n$, respectively. Well,

$$\begin{aligned} \text{Suppose } & d = b_1 \cup \dots \cup b_n \\ \text{So } & \mathcal{C} \models \text{Sum}(b_1, \dots, b_n, d) \\ \text{So } & \mathcal{S}_{\mathcal{C}} \models \text{Sum}(\sigma_{\mathcal{B}}[b_1], \dots, \sigma_{\mathcal{B}}[b_n], \sigma_{\mathcal{B}}[d]) \\ \text{So } & \mathcal{S}_{\mathcal{A}} \models \text{Sum}(g(\sigma_{\mathcal{B}}[b_1]), \dots, g(\sigma_{\mathcal{B}}[b_n]), g(\sigma_{\mathcal{B}}[d])) \\ \text{So } & \mathcal{S}_{\mathcal{A}} \models \text{Sum}(c_1, \dots, c_n, d) \end{aligned}$$

since each $c_i \in g(\sigma_{\mathcal{B}}[b_i])$ and $d \in g(\sigma_{\mathcal{B}}[d]) = \sigma_{\mathcal{A}}[d]$. So, the existence of $a_1, \dots, a_n$, as above, is guaranteed because $\mathcal{A} \models \text{DEF}^+$.

Now, let $f(b_i) = a_i$ for each b_i in the molecule d . Repeating this procedure for each molecule yields a value of f for each atom of $\mathcal{C}$. Finally, if $x \in \dot{\mathcal{C}}$ is non-atomic, then

$$x = b_1 \cup \dots \cup b_k$$

where each b_k is atomic. So let

$$f(x) = f(b_1) \cup \dots \cup f(b_k)$$

f satisfies the requirements of Theorem 5.1.2: Boolean relations are preserved by f because the function is determined by its values on the atoms of $\mathcal{C}$; f maps elements of $\mathcal{C} \cap A$ into themselves because the set of atoms contained in each of these molecules is mapped into a disjoint collection of elements of $\dot{A}$ whose union is the same molecule; so, we need only show that f preserves size relations. To do so, we invoke lemma 5.1.6, below:

- (a) $\mathcal{C} \models x < y$ iff $\mathcal{A} \models f(x) < f(y)$
- (b) $\mathcal{C} \models x \sim y$ iff $\mathcal{A} \models f(x) \sim f(y)$
- (c) $\mathcal{C} \models \text{Sum}(x, y, z)$ iff $\mathcal{A} \models \text{Sum}(f(x), f(y), f(z))$
- (d) $\mathcal{C} \models \text{Unit}(x)$ iff $\mathcal{A} \models \text{Unit}(y)$

Proof of (a): (The others are similar).

$$\begin{aligned}
 \mathcal{C} \models x < y &\text{ iff } \mathcal{S}_{\mathcal{B}} \models \sigma_{\mathcal{B}}[x] < \sigma_{\mathcal{B}}[y] \\
 &\text{ iff } \mathcal{S}_{\mathcal{A}} \models g(\sigma_{\mathcal{B}}[x]) < g(\sigma_{\mathcal{B}}[y]) \\
 &\text{ iff } \mathcal{S}_{\mathcal{A}} \models \sigma_{\mathcal{A}}[f(x)] < \sigma_{\mathcal{A}}[f(y)] && \text{by 5.1.6} \\
 &\text{ iff } \mathcal{A} \models f(x) < f(y)
 \end{aligned}$$

□

So, we may conclude:

Theorem 5.1.5. *CA is model complete.*

Lemma 5.1.6. *For all x in $\dot{\mathcal{C}}$,*

$$\sigma_{\mathcal{A}}[f(x)] = g(\sigma_{\mathcal{B}}[x])$$

Proof. Suppose

$$x = b_1 \cup \dots \cup b_n$$

where each b_i is an atom of $\dot{\mathcal{C}}$.

$$\begin{aligned}
 \text{So } \mathcal{B} \models \text{Sum}(b_1, \dots, b_n, x) &&& \text{since } \mathcal{B} \models \text{DISJ}^\cup \\
 \text{So } \mathcal{S}_{\mathcal{B}} \models \text{Sum}(\sigma_{\mathcal{B}}[b_1], \dots, \sigma_{\mathcal{B}}[b_n], \sigma_{\mathcal{B}}[x]) &&& \\
 \text{So } \mathcal{S}_{\mathcal{A}} \models \text{Sum}(g(\sigma_{\mathcal{B}}[b_1]), \dots, g(\sigma_{\mathcal{B}}[b_n]), g(\sigma_{\mathcal{B}}[x])) &&& \text{since } \mathcal{S}_{\mathcal{A}} \subseteq \mathcal{S}_{\mathcal{B}} \\
 \text{But } g(\sigma_{\mathcal{B}}[b]) = \sigma_{\mathcal{A}}[f(b)] &&& \text{by the choice of } f(b). \\
 \text{So } \mathcal{S}_{\mathcal{A}} \models \text{Sum}(\sigma_{\mathcal{A}}[f(b_1)], \dots, \sigma_{\mathcal{A}}[f(b_n)], g(\sigma_{\mathcal{B}}[x])) &&& \\
 \text{But } \mathcal{A} \models \text{Sum}(f(b_1), \dots, f(b_n), f(x)) &&& \text{since } \mathcal{A} \models \text{DISJ}^\cup \\
 \text{So } \mathcal{S}_{\mathcal{A}} \models \text{Sum}(\sigma_{\mathcal{A}}[f(b_1)], \dots, \sigma_{\mathcal{A}}[f(b_n)], \sigma_{\mathcal{A}}[f(x)]) &&& \\
 \text{So } g(\sigma_{\mathcal{B}}[x]) = \sigma_{\mathcal{A}}[f(x)] &&& \text{since } \mathcal{S}_{\mathcal{A}} \models \text{UNIQ}^+
 \end{aligned}$$

□

5.2 Prime models for CA

For each total, congruous remainder function, f , we want to find a prime model, $\mathcal{Q}_f$, for CAI_f . All of these prime models can be defined over the class, $\mathcal{Q}$, of sets near quasi-congruence classes (see section 3.2). For different remainder functions, we need to assign different size relations over $\mathcal{Q}$. Section 5.2.1 defines the structures $\mathcal{Q}_f$ and verifies that each satisfies the respective theory, CAI_f . Section 5.2.2 defines, for each model of CAI_f , a submodel, or *shell*. Section 5.2.3 shows that $\mathcal{Q}_f$ is isomorphic to the shell of any model of CAI_f .

5.2.1 The standard protomodels

The construction here is a more elaborate version of the construction of $\mathcal{Q}$ in chapter 3 (see 3.2.19). $\mathcal{Q}$ turns out to be $\mathcal{Q}_f$, where $f(n) = 0$ for all n . As in the case of $\mathcal{Q}$, the models $\mathcal{Q}_f$ and their copies in arbitrary models of CAI are unions of chains.

The sizes assigned to elements of $\mathcal{Q}$ to induce $\mathcal{Q}_f$ for a total, congruous remainder function, f , are more elaborate than those used in the definition of $\mathcal{Q}$ (see 3.2.22), but they are employed in substantially the same way:

Definition 5.2.1.

*a. A **size** is an ordered pair $\langle \rho, \delta \rangle$, where both ρ and δ are rational.*

b. If $\theta_1 = \langle \rho_1, \delta_1 \rangle$ and $\theta_2 = \langle \rho_2, \delta_2 \rangle$ are sizes, then

(i) $\theta_1 < \theta_2$ iff $\rho_1 < \rho_2$ or $\rho_1 = \rho_2$ and $\delta_1 < \delta_2$.

(ii) $\theta_1 + \theta_2 = \langle \rho_1 + \rho_2, \delta_1 + \delta_2 \rangle$

(cf 3.2.22)

To assign sizes for $\mathcal{Q}_f$ we rely on the representation of sets in $\mathcal{Q}$ defined in 3.2.24. $\mathcal{Q}_f$, unlike $\mathcal{Q}$, assigns different sizes to the n -congruence classes for a given n .

Definition 5.2.2. *If f is total and congruous, then*

a. If x is an n -congruence class, $x = [n\mathbf{k} + i]$, then

$$\theta_f(x) = \begin{cases} \langle \frac{1}{n}, \frac{n-f(n)}{n} \rangle, & \text{if } i < f(n), \\ \langle \frac{1}{n}, \frac{-f(n)}{n} \rangle, & \text{if } f(n) \leq i. \end{cases}$$

b. If $x \in \mathcal{QC}_n$, so that x is the disjoint union of n -congruence classes,

$$x_1 \cup \dots \cup x_k$$

then

$$\theta_f(x) = \theta_f(x_1) + \dots + \theta_f(x_k)$$

c. If x is finite, then

$$\theta_f(x) = \langle 0, |x| \rangle$$

d. If $x \in Q$, so that x can be represented as

$$(C(x) \cup \Delta_1(x)) - \Delta_2(x)$$

as in 3.2.24, then

$$\begin{aligned} \theta_f(x) &= \theta_f(C(x)) + \theta_f(\Delta_1(x)) - \theta_f(\Delta_2(x)) \\ &= \theta_f(C(x)) + \langle 0, |\Delta_1(x)| \rangle - \langle 0, |\Delta_2(x)| \rangle \\ &= \theta_f(C(x)) + \langle 0, |\Delta_1(x)| - |\Delta_2(x)| \rangle \end{aligned}$$

Intuitively, all n -congruence classes are assigned sizes $\langle 1/n, \delta \rangle$, but δ is no longer 0 in all cases, as in Q . Instead, the first $f(n)$ n -congruence classes are each one atom larger than the remaining $(n - f(n))$ n -congruence classes.

The desired models of CAI may now be defined:

Definition 5.2.3. If f is total and congruous, then the **standard protomodel** for f is Q_f , where

$$\begin{aligned} \dot{Q}_f &= Q \\ Q_f &\models x < y \text{ iff } \theta_f(x) < \theta_f(y) \\ Q_f &\models x \sim y \text{ iff } \theta_f(x) = \theta_f(y) \\ Q_f &\models \text{Unit}(x) \text{ iff } \theta_f(x) = \langle 0, 1 \rangle \\ Q_f &\models \text{Sum}(x, y, z) \text{ iff } \theta_f(z) = \theta_f(x) + \theta_f(y) \end{aligned}$$

(cf 3.2.27)

To verify that the structure Q_f is a model of CAI_f , for total and congruous f , we exhibit each such model as the union of a chain of models.

Definition 5.2.4. If f is total and congruous, then $Q_{f,n}$ is the submodel of Q_f whose domain is Q_n .

Fact 5.2.5. If f is total and congruous and $n > 0$, then $Q_{f,n} \models \text{BASIC}$.

Proof. The proof can be obtained from the proof of Theorem 3.2.29 by substituting:

$$\begin{aligned} Q_{f,n} &\text{ for } Q_n, \\ \theta_f(x) &\text{ for } \theta(x), \\ \rho_f(x) &\text{ for } \rho(x), \text{ and} \\ \delta_f(x) &\text{ for } \delta(x) \end{aligned}$$

□

The following notion is helpful in understanding our constructions.

Definition 5.2.6. Suppose $\mathcal{A} \models \text{BASIC}$, $x \in \dot{\mathcal{A}}$, and $0 \leq m < n$. Then an $(n.m)$ -**partition** of x in $\mathcal{A}$ is a sequence $x_1, \dots, x_n$, where

- a. $x = x_1 \cup \dots \cup x_n$
- b. If $0 < i < j \leq n$, then $x_i \cap x_j = \emptyset$, and
- c. The x_i are approximately the same size:
 - (i) If $0 < i < j \leq m$, then $x_i \sim x_j$,
 - (ii) If $m < i < j \leq n$, then $x_i \sim x_j$,
 - (iii) If $0 < i \leq m < j \leq n$, then $\text{Sum}(x_i, a, x_j)$, for any atom a ,

In other words, x is partitioned among n pairwise disjoint sets which are roughly the same size: each of the first m is one atom larger than each of the remaining $n - m$. If $0 < i \leq m$, x_i is called a **charmed n -factor** of x ; for $i > m$, x_i is a **common n -factor** of x .

The sequence $\mathcal{Q}_{f,n}$ does not constitute a chain of models. For example $\mathcal{Q}_{f,3}$ is not an extension of $\mathcal{Q}_{f,2}$. But this sequence does harbor a chain of models:

Fact 5.2.7. If $n > m$, then $\mathcal{Q}_{f,n!} \subseteq \mathcal{Q}_{f,m!}$.

Proof. It will be clearer, and easier, to establish this by example rather than by formal proof. Letting $n = 2$ and $m = 3$, we want to show that the 2-congruence classes have the same size relations in $\mathcal{Q}_{f,6}$ as they do in $\mathcal{Q}_{f,2}$, for any f . The other elements of $\mathcal{Q}_{f,6}$ will then fall into place, since size relations are determined by the representations of each set, x , as $C(x)$, $\Delta_1(x)$, and $\Delta_2(x)$.

Suppose that $f(2) = 0$, so that

$$\mathcal{Q}_{f,2} \models [2\mathbf{n}] \sim [2\mathbf{n} + 1]$$

Since f is congruous, $f(6) \in \{0, 2, 4\}$. If $f(6) = 0$, then all of the 6-congruence classes are common. If $f(6) = 2$, then $[6\mathbf{n}]$ and $[6\mathbf{n} + 1]$ are the only charmed 3-congruence classes. If $f(6) = 4$, then all of the 3-congruence classes are charmed except for $[6\mathbf{n} + 4]$ and $[6\mathbf{n} + 5]$.

In any case, $[2\mathbf{n}]$ will include the same number of charmed 3-congruence classes as $[2\mathbf{n} + 1]$, so

$$\mathcal{Q}_{f,6} \models [2\mathbf{n}] \sim [2\mathbf{n} + 1]$$

Suppose, however that $f(2) = 1$, so that

$$\mathcal{Q}_{f,2} \models [2\mathbf{n}] \text{ is one atom larger than } [2\mathbf{n} + 1]$$

Here, $f(6) \in \{1, 3, 5\}$, since f is congruous. In any case, $[2\mathbf{n}]$ contains exactly one more charmed congruence class than $[2\mathbf{n} + 1]$, so

$$\mathcal{Q}_{f,6} \models [2\mathbf{n}] \text{ is one atom larger than } [2\mathbf{n} + 1]$$

So it goes in general. □

Fact 5.2.7 allows us to regard $\mathcal{Q}_f$ as the union of a chain of models:

Fact 5.2.8. *If f is total and congruous, then*

$$\mathcal{Q}_f = \bigcup_{n>0} \mathcal{Q}_{f,n!}$$

Fact 5.2.9. *If f is total and congruous, then*

- a. $\mathcal{Q}_f \models \text{BASIC}$
- b. $\mathcal{Q}_f \models \text{ADIV}_k$, for $k > 0$
- c. $\mathcal{Q}_f \models \text{MOD}_{n,f(n)}$, for $n > 0$
- d. $\mathcal{Q}_f \models \text{CAI}_f$

Proof.

- a. BASIC is a universal-existential theory; so it is preserved under unions of chains. (See Appendix, Fact B.5).
- b. Each n -congruence class can be partitioned into k nk -congruence classes.
- c. $\mathcal{Q}_{f,n} \models \text{MOD}_{n,f(n)}$, by definition. Since $\text{MOD}_{n,f(n)}$ is an existential sentence, it is preserved under extensions.
- d. Immediate from (a), (b), and (c).

□

5.2.2 Shells of models

To embed the model $\mathcal{Q}_f$ into an arbitrary model $\mathcal{A}$, of CAI_f , we must find a smallest submodel $\mathcal{B}$, of $\mathcal{A}$ which satisfies CAI. Clearly, the basis of $\mathcal{A}$, call it x_0 , must be included in $\mathcal{B}$, since the symbol **I** must refer to the same set in the submodel as it does in the model. But, if the basis of $\mathcal{A}$ is in $\dot{\mathcal{B}}$ and $\mathcal{B} \models \text{CAI}$, then $\dot{\mathcal{B}}$ must contain two disjoint sets of roughly the same size whose union is the basis of $\mathcal{A}$, x_0 . Pick such a pair, x_1 and x_2 , to include in $\dot{\mathcal{B}}$. Whether these are exactly the same size or differ by an atom depends on whether $\mathcal{A} \models \text{MOD}_{2,0}$ or $\mathcal{A} \models \text{MOD}_{2,1}$.

$\mathcal{B}$ must also satisfy ADIV_3 . We can aim for this by placing in $\dot{\mathcal{B}}$ three disjoint sets, x_{11} , x_{12} , and x_{13} whose union is x_1 and another three disjoint sets, x_{21} , x_{22} , and x_{23} whose union is x_2 . The existence of such sets is assured because $\mathcal{A} \models \text{ADIV}_3$. Again, the exact size relations will be determined by which MOD principles are satisfied in $\mathcal{A}$. Insuring that x_1 and x_2 are each divisible by 3, also guarantees that x_0 is divisible by 3: the three unions

$$x_{11} \cup x_{21}, x_{12} \cup x_{22}, \text{ and } x_{13} \cup x_{23}$$

will be roughly the same size and will exhaust x_0 .

We can continue this process indefinitely, dividing each set introduced at stage n into $n + 1$ roughly equal subsets at stage $n + 1$. This will produce an infinite tree, bearing sets. The deeper a node is in this tree, the smaller the set it bears and the greater the number of successors among which this set will be partitioned.

This great tree of sets will not form a boolean algebra, nor will it be closed under finite unions. A boolean algebra could be obtained by including both the node sets and their finite unions, but this would still not be an atomic boolean algebra, which is what we are looking for. We cannot correct for this problem by including in $\tilde{\mathcal{B}}$ all atoms of $\mathcal{A}$: $\mathcal{A}$ may have uncountably many atoms while $\mathcal{B}$, to be a prime model, must be countable. We leave the solution of this problem to the formal construction.

The formal proof proceeds as follows: First, we define a tree, i.e. the set of nodes on which we shall hang both the components of the successive partitions described above and the atoms of the submodel being constructed. Second, we present the construction which, given a model $\mathcal{A}$ of CAI_f , assigns a **node set** $\mathbf{A}_\lambda$ and a **node atom**, $\mathbf{a}_\lambda$, to each node, λ . Third, we define the **shell** of $\mathcal{A}$ as the submodel of $\mathcal{A}$ generated by the collection of node sets and node atoms. In the next section, we show that the shell of $\mathcal{A}$ is isomorphic to $\mathcal{Q}_f$.

First, the tree:

Definition 5.2.10.

- a. A **node** is a finite sequence $\langle n_1, \dots, n_k \rangle$, where $k > 0$ and for all $i \leq k$, $n_i < i$. (The variables λ and κ range over nodes.)
- b. If $\lambda = \langle n_1, \dots, n_k \rangle$, then
 - (i) The **length**, or **depth**, of λ , $\mathbf{L}(\lambda)$ is k .
 - (ii) If $1 \leq i \leq k$, then $\lambda(i) = n_i$.
 - (iii) $\lambda \bullet m = \langle n_1, \dots, n_k, m \rangle$
- c. λ **extends** κ iff $\mathbf{L}(\lambda) > \mathbf{L}(\kappa)$ and, for $1 \leq i \leq \mathbf{L}(\kappa)$, $\lambda(i) = \kappa(i)$.
- d. λ **0-extends** κ iff λ extends κ and, for $\mathbf{L}(\kappa) < i \leq \mathbf{L}(\lambda)$, $\lambda(i) = 0$.

Nodes constitute the vertices of an infinite tree in which $\langle 0 \rangle$ is the root and λ dominates κ iff κ extends λ . The number of immediate descendants of a node grows as the depth of the node increases.

We shall now assign a set to each node by repeatedly partitioning the basis of $\mathcal{A}$. At the same time we shall assign an atom to each node.

Definition 5.2.11. Given $\mathcal{A} = \mathcal{A}_f \models CAI_f$, where f is a total, congruous remainder function:

- a. Let $\mathbf{A}_{\langle 0 \rangle}$ be the basis of $\mathcal{A}$, and let $\mathbf{a}_{\langle 0 \rangle}$ be any atom of $\mathcal{A}$.

b. Suppose that $\mathbf{A}_\lambda$ and $\mathbf{a}_\lambda$ have been chosen. Let $m = \mathbf{L}(\lambda)$ and let

$$k = \frac{f((m+1)!) - f(m!)}{m!}$$

Since f is congruous, k is an integer (see Definition 3.3.7e). Let

$$\mathbf{A}_{\lambda \bullet 0}, \dots, \mathbf{A}_{\lambda \bullet m}$$

be an $(m+1, k)$ partition of $\mathbf{A}_\lambda$ if $\mathbf{A}_\lambda$ is common or an $(m+1, k+1)$ partition of $\mathbf{A}_\lambda$ if $\mathbf{A}_\lambda$ is charmed. Fact 5.2.12 guarantees that such partitions exist. In either case, choose $\mathbf{A}_{\lambda \bullet 0}$ so that it contains $\mathbf{a}_\lambda$. This is always possible because $\mathbf{A}_\lambda$ contains $\mathbf{a}_\lambda$.

c. Let $\mathbf{a}_{\lambda \bullet 0} = \mathbf{a}_\lambda$. If $0 < i \leq m$, let $\mathbf{a}_{\lambda \bullet i}$ be any atomic subset of $\mathbf{A}_{\lambda \bullet i}$.

The sets $\mathbf{A}_\lambda$ will be referred to as **node sets** and the atoms $\mathbf{a}_\lambda$ as **node atoms**.

Fact 5.2.12. Suppose $\mathcal{A} \models \text{CAI}_f$, $n > 0$, $m > 0$, and

$$k = \frac{f(nm) - f(n)}{n}$$

Then

- a. x has an (m, k) -partition iff $\mathcal{A} \models \text{Mod}_{m,k}(x)$
- b. Every common n -factor of $\dot{\mathcal{A}}$ has an (m, k) -partition.
- c. Every charmed n -factor of $\dot{\mathcal{A}}$ has an $(m, k+1)$ -partition.

Proof.

- a. $\text{Mod}_{m,k}(x)$ says that x has an (m, k) -partition.
- b. All common n -factors are the same size and satisfy the same $\text{Mod}_{m,k}$ predicate. So, suppose that each common n -factor has k charmed m -factors and $m - k$ common m -factors.

By (a), $\mathcal{A}$ has $f(n)$ charmed n -factors and $n - f(n)$ common n -factors. Partitioning each of the n -factors into m subsets of roughly the same size yields an nm -partition of $\mathcal{A}$; the charmed m -factors of the n -factors are the charmed nm factors of $\mathcal{A}$ and the common m -factors of the n -factors are the common nm factors of $\mathcal{A}$.

Each of the common n factors has k charmed m -factors and each of the charmed n -factors has $k+1$ charmed m -factors. In all, there are

$$\begin{aligned} & (n - f(n))k + f(n)(k+1) \\ &= nk + f(n) \end{aligned}$$

charmed m -factors among the n -factors of $\mathcal{A}$.

But, by (a) again, there are $f(nm)$ charmed nm -factors of $\mathcal{A}$. So

$$\begin{aligned} f(nm) &= nk + f(n) \\ \text{and } k &= (f(nm) - f(n))/n \end{aligned}$$

- c. Immediate from (b), since every charmed n -factor is one atom larger than each common n -factor.

□

The facts listed in 5.2.13 should clarify this constuction. All of them can be established by induction on the depth of nodes.

Fact 5.2.13.

- a. $\mathbf{A}_\lambda \subset \mathbf{A}_\kappa$ iff κ extends λ or $\kappa = \lambda$.
- b. If $\mathbf{A}_\lambda = \mathbf{A}_\kappa$, then $\kappa = \lambda$.
- c. There are $n!$ nodes (and, hence, node sets) of depth n .
- d. If $i \neq j$, then $\mathbf{A}_{\lambda \bullet i} \cap \mathbf{A}_{\lambda \bullet j} = \emptyset$.
- e. Any two node sets of the same depth are disjoint.
- f. Each node set is the disjoint union of its immediate descendants.
- g. Each node set is the disjoint union of all of its descendants at any given depth.
- h. $\mathbf{a}_\lambda \subset \mathbf{A}_\kappa$ iff λ extends κ .
- i. $\mathbf{a}_\lambda = \mathbf{a}_\kappa$ iff one of λ and κ 0-extends the other.
- j. Every node set contains infinitely many node atoms.
- k. For any n , the node sets of depth n form an $(n!, f(n!))$ -partition of the basis of $\mathcal{A}$.

Proof. We demonstrate (k) by induction on n :. If $n = 1$, then $n! = 1$, $f(n!) = 0$, and $\mathbf{A}_{\lambda(0)}$ is the basis of $\mathcal{A}$. So (k) holds because any set is a $(1,0)$ -partition of itself.

Assume that (k) holds for n . Then (k) also holds for $n + 1$: each node set of depth n has $n + 1$ immediate descendants; so, there are $(n!)(n + 1) = (n + 1)!$ node sets of depth $n + 1$.

Furthermore, $f(n!)$ of the n -factors are charmed and $n! - f(n!)$ are common. By Fact 5.2.12, each charmed n -factor has an $(n + 1, k + 1)$ -partition and each common n -factor has an $(n + 1, k)$ -partition, where

$$\begin{aligned} k &= \frac{f(n!(n + 1)) - f(n!)}{n!} \\ &= \frac{f((n + 1)!) - f(n!)}{n!} \end{aligned}$$

(substituting $n!$ for n and $n + 1$ for m).

So, there are

$(k + 1)f(n!)$ charmed $n + 1$ factors from the charmed n -factors
and $k(n! - f(n!))$ charmed $n + 1$ factors from the common n -factors

In all, then, the number of charmed n -factors is:

$$\begin{aligned}
 & (k + 1)f(n!) + k(n! - f(n!)) \\
 &= kf(n!) + f(n!) + n!k - f(n!)k \\
 &= f(n!) + n!k \\
 &= f(n!) + f((n!)(n + 1)) - f(n!) \\
 &= f(n!)(n + 1) \\
 &= f((n + 1)!)
 \end{aligned}$$

So, the $n + 1$ factors of the n -factors of the basis form an $((n + 1)!, f((n + 1)!))$ -partition of the basis. That is to say, (k) holds for $n + 1$. $\square$

Given a collection of node sets, $\mathbf{A}_\lambda$, and node-atoms, $\mathbf{a}_\lambda$, from a model, $\mathcal{A}$, of CAI_f , we can now construct a submodel, $\mathcal{B}$, of $\mathcal{A}$ which is isomorphic to $\mathcal{Q}_f$.

Definition 5.2.14. Suppose f is total and congruous, that $\mathcal{A} \models \text{CAI}_f$ and $\mathbf{A}_\lambda$ and $\mathbf{a}_\lambda$ are the node sets and node-atoms of $\mathcal{A}$ produced by construction 5.1.12. Then, the *shell* of $\mathcal{A}$ is the submodel of $\mathcal{A}$ generated by $\{\mathbf{A}_\lambda, \mathbf{a}_\lambda\}$.

For the remainder of this chapter, we will regard as fixed:

- f , a total, congruous remainder function
- $\mathcal{A}$, a model of CAI_f ,
- $\{\mathbf{A}_\lambda, \mathbf{a}_\lambda\}$, a set of node sets and node-atoms produced by the construction above.
- $\hat{\mathcal{A}}$, the shell of $\mathcal{A}$ generated from $\{\mathbf{A}_\lambda, \mathbf{a}_\lambda\}$

To show that $\hat{\mathcal{A}} \simeq \mathcal{Q}_f$, we need a sharper characterization of the elements of $\hat{\mathcal{A}}$. Recall from 3.2.24 that each member, x , of $\mathcal{Q}$ has a unique representation as

$$(C(x) \cup \Delta_1(x)) - \Delta_2(x)$$

where $C(x)$ is a quasi-congruence class, $\Delta_1(x)$ and $\Delta_2(x)$ are finite sets and

$$\begin{aligned}
 \Delta_1(x) \cap C(x) &= \emptyset \\
 \Delta_1(x) \cap \Delta_2(x) &= \emptyset \\
 \Delta_2(x) &\subseteq C(x)
 \end{aligned}$$

We can obtain a similar representation for elements of $\hat{\mathcal{A}}$: the node sets play the role of (some of) the congruence classes; finite unions of node sets correspond to the quasi-congruence classes; finite sets of node-atoms correspond to the finite subsets in Q .

Definition 5.2.15.

- a. x is a **quasi-nodal** set of $\mathcal{A}$ iff it is the union of finitely many node sets (iff it is the union of finitely many node sets at a given depth).
- b. x is an **$\mathcal{A}$ -finite set** iff it is a finite set of node atoms.
- c. If $x \in A$ and $y \in A$, then x is **$\mathcal{A}$ -near** y iff both $x - y$ and $y - x$ are $\mathcal{A}$ -finite sets,

(cf. 3.2.14-3.2.18)

Still following in the footsteps of chapter 3, we can characterize $\hat{\mathcal{A}}$ as the collection of sets $\mathcal{A}$ -near quasi-nodal sets. Analogues of 3.2.15 through 3.2.18 obtain for $\mathcal{A}$ -nearness.

Fact 5.2.16.

- a. $x \in \hat{\mathcal{A}}$ iff x is $\mathcal{A}$ -near some quasi-nodal set of $\mathcal{A}$.
- b. $\hat{\mathcal{A}}$ is an atomic boolean algebra whose atoms are the node-atoms $\mathbf{a}_\lambda$.
- c. If $x \in \hat{\mathcal{A}}$, then x has a unique representation as

$$(C(x) \cup \Delta_1(x)) - \Delta_2(x)$$

where $C(x)$ is a quasi-nodal set disjoint from $\Delta_1(x)$ and including $\Delta_2(x)$, both of which are $\mathcal{A}$ -finite sets.

Proof.

- a. $(\Rightarrow)$ $\hat{\mathcal{A}}$ is generated from node sets and node atoms via the boolean operations, each of which preserves $\mathcal{A}$ -nearness to quasi-nodal sets.
 $(\Leftarrow)$ $\hat{\mathcal{A}}$ must contain finite unions of node sets as well as $\mathcal{A}$ -finite sets; so it must also contain sets obtained by adding or removing $\mathcal{A}$ -finite sets from quasi-nodal sets.
- b. The proof parallels that of Theorem 3.2.21 exactly.
- c. Let $C(x)$ be the quasi-nodal set which is $\mathcal{A}$ -near x (see 3.2.23); Let $\Delta_1(x) = x - C(x)$; and, let $\Delta_2(x) = C(x) - x$.

□

5.2.3 The embeddings

To embed $\mathcal{Q}_f$ into $\mathcal{A}$, we first describe Q in terms of node sets and node atoms. In effect, we are performing the construction 5.2.12 on $\mathcal{Q}_f$, but with two differences: first, we are stipulating which (n, m) -partitions to use at each level; second, we are selecting node atoms so that every singleton in Q is the node atom for some node. This latter condition guarantees that the shell of $\mathcal{Q}_f$ will be $\mathcal{Q}_f$ itself.

Definition 5.2.17. *Suppose λ is a node. Then*

- a. *If $\mathbf{L}(\lambda) = k$, then*

$$Q_\lambda = [k! \mathbf{n} + m]$$

where

$$m = \sum_{i=1}^k \lambda(i)(i-1)!$$

- b. *The **depth** of Q_λ is $\mathbf{L}(\lambda)$.*

Examples.

$$\begin{aligned} \mathbf{Q}_{\langle 0 \rangle} &= [\mathbf{n}] \\ \mathbf{Q}_{\langle 0,0 \rangle} &= [2\mathbf{n}] \\ \mathbf{Q}_{\langle 0,1 \rangle} &= [2\mathbf{n} + 1] \\ \mathbf{Q}_{\langle 0,1,0 \rangle} &= [6\mathbf{n} + 1] \\ \mathbf{Q}_{\langle 0,0,2 \rangle} &= [6\mathbf{n} + 4] \\ \mathbf{Q}_{\langle 0,1,2 \rangle} &= [6\mathbf{n} + 5] \end{aligned}$$

Definition 5.2.18.

- a. $\iota(\lambda) =$ *the least $n \in Q_\lambda$.*
b. $\mathbf{q}_\lambda = \{\iota(\lambda)\}$

Fact 5.2.19.

- a. *If $\lambda = \langle n_1, \dots, n_k \rangle$, then*

$$\iota(\lambda) = \sum_{i=1}^k (i-1)!n_i$$

- b. $\iota(\lambda) = \iota(\kappa)$ *iff $\lambda = \kappa$ or one 0-extends the other.*

- c. For every n , there's a λ such that $n = \iota(\lambda)$.
- d. For every n , there are infinitely many λ such that $n = \iota(\lambda)$.
- e. For every n , there's a λ such that $n = \iota(\kappa)$ iff $\kappa = \lambda$ or κ 0-extends λ .

Fact 5.2.20.

- a. At each depth, n , the $\iota(\lambda)$ take on all and only values less than $n!$.
- b. If $\iota(\lambda) = k$, then all nodes along the left-most branch descending from λ also have the value k . These are the only nodes below λ with the value k .
- c. Every natural number is the value of all and only those nodes along the left-most branch descending from some node.

Though for a given natural number n , there will be infinitely many nodes λ for which $\iota(\lambda) = n$, we can associate with each natural number a shortest (i.e. shallowest) node for which $\iota(\lambda) = n$.

Definition 5.2.21. λ_n = the shortest λ such that $\iota(\lambda) = n$.

Fact 5.2.22.

- a. $\iota(\lambda_n) = n$
- b. λ extends $\lambda_{\iota(\lambda)}$
- c. $\lambda_{\iota(\lambda_n)} = \lambda_n$
- d. $\lambda_{\iota(\lambda)} = \lambda$ iff $\lambda = \langle 0 \rangle$ or $\lambda(\mathbf{L}(\lambda)) \neq 0$ (ie a node, λ , will be the highest node with a certain value just in case λ is not the leftmost immediate descendant of its parent.)

Each of the points listed in Fact 5.2.13 hold for the sets Q_λ and $\mathbf{q}_\lambda$. That is to say, the Q_λ can be regarded as node sets and the $\mathbf{q}_\lambda$ as node atoms for any model $\mathcal{Q}_f$. Notice, especially, that 5.2.13k holds.

We may, finally, define the embedding of $\mathcal{Q}_f$ into $\mathcal{A}$:

Fact 5.2.23. If $x \in Q$, there is a unique $y \in \dot{\mathcal{A}}$ such that

$$\forall \lambda (\mathbf{q}_\lambda \subseteq x \leftrightarrow \mathbf{a}_\lambda \subseteq y)$$

Proof. There is at most one such y , by Fact 5.2.16. To show that there is such a y , suppose first that $x \in QC$.

Then there is some n such that

$$x = x_1 \cup \dots \cup x_k$$

where each x_i is an $(n!)$ -congruence class and hence a node set in $\mathcal{Q}_f$. So

$$x = \mathbf{Q}_{\lambda_1} \cup \cdots \cup \mathbf{Q}_{\lambda_k}$$

Now, let

$$y = \mathbf{A}_{\lambda_1} \cup \cdots \cup \mathbf{A}_{\lambda_k}$$

So $y \in \dot{\mathcal{A}}$ and:

$$\begin{aligned} \mathbf{q}_\lambda \subset x &\text{ iff } \mathbf{q}_\lambda \subset \mathbf{Q}_{\lambda_i}, \text{ for some } i \\ &\text{ iff } \lambda \text{ extends } \lambda_i && \text{ by 5.2..13h} \\ &\text{ iff } \mathbf{a}_\lambda \subset \mathbf{A}_{\lambda_i} \\ &\text{ iff } \mathbf{a}_\lambda \subset y. \end{aligned}$$

If x is not a quasi-congruence class, then

$$x = (x' \cup \Delta_1(x)) - \Delta_2(x)$$

where x' is a quasi-congruence class. Let y' be the element of $\dot{\mathcal{A}}$ corresponding to x' , as described above, and let

$$y = (y' \cup \{ \mathbf{a}_\lambda \mid \mathbf{q}_\lambda \subseteq \Delta_1(x) \}) - \{ \mathbf{a}_\lambda \mid \mathbf{q}_\lambda \subseteq \Delta_2(x) \}$$

□

Definition 5.2.24. The *nodal embedding* of Q into $\dot{\mathcal{A}}$ is defined by letting $g(x \in Q)$ be the $y \in \dot{\mathcal{A}}$ such that

$$\forall \lambda (\mathbf{a}_\lambda \subseteq y \leftrightarrow \mathbf{q}_\lambda \subseteq x)$$

Fact 5.2.25.

a. If $x \in Q$, then

$$g(x) = (g(C(x)) \cup g(\Delta_1(x))) - g(\Delta_2(x))$$

b. g is one-one.

c. g maps Q onto $\dot{\mathcal{A}}$.

Proof.

a. Immediate from the proof of 5.2.23.

b. Suppose $g(x) = y = g(x')$. Then

$$\forall \lambda (\mathbf{q}_\lambda \subseteq x \leftrightarrow \mathbf{a}_\lambda \subseteq y \leftrightarrow \mathbf{q}_\lambda \subseteq x')$$

But every integer is $\iota(\lambda)$ for some λ , so $x = x'$.

c. Obvious.

□

Theorem 5.2.26. *The nodal embedding, g , of Q into $\mathcal{A}$ is an isomorphism of $\mathcal{Q}_f$ onto $\hat{\mathcal{A}}$.*

Proof.

a.

$$\begin{aligned} \mathcal{Q}_f \models x \subseteq y &\text{ iff } \forall \lambda (\mathbf{q}_\lambda \subset x \rightarrow \mathbf{q}_\lambda \subset y) \\ &\text{ iff } \forall \lambda (\mathbf{a}_\lambda \subseteq g(x) \rightarrow \mathbf{a}_\lambda \subseteq g(y)) && \text{(by 5.2.24)} \\ &\text{ iff } \hat{\mathcal{A}} \models g(x) \subseteq g(y) && \text{(by 5.2.16)} \end{aligned}$$

b. As in (a), g can be shown to preserve $\emptyset$, $\mathbf{I}$, unions, intersections, relative complements, and proper subsets.

c. Let $\text{nc}(z, n)$ be the number of charmed node sets of level n contained in $C(z)$. And let n , below, be the least k such that x and y are both unions of node sets of depth k .

$$\begin{aligned} \mathcal{Q}_f \models x \sim y &\text{ iff } \theta_f(x) = \theta_f(y) \\ &\text{ iff } \text{nc}(x, n) - (|\Delta_1(x)| - |\Delta_2(x)|) \\ &\quad = \text{nc}(y, n) - (|\Delta_1(y)| - |\Delta_2(y)|) \\ &\text{ iff } \text{nc}(g(x), n) - (|g(\Delta_1(x))| - |g(\Delta_2(x))|) \\ &\quad = \text{nc}(g(y), n) - (|g(\Delta_1(y))| - |g(\Delta_2(y))|) \\ &\quad \text{since } Q_\lambda \text{ is charmed iff } \mathbf{A}_\lambda \text{ is charmed} \\ &\quad \text{and } g \text{ preserves boolean relations} \\ &\text{ iff } \mathcal{A} \models g(x) \sim g(y) \\ &\text{ iff } \hat{\mathcal{A}} \models g(x) \sim g(y) && \text{since } \hat{\mathcal{A}} \subseteq \mathcal{A} \end{aligned}$$

d.

$$\begin{aligned} \mathcal{Q}_f \models x < y &\text{ iff } \mathcal{Q}_f \models x \sim x' \subset y \\ &\quad \text{for some } x' \in Q && \text{since } \mathcal{Q}_f \models \text{REP}^< \\ &\text{ iff } \mathcal{A} \models g(x) \sim g(x') \subset g(y) && \text{by (b) and (c)} \\ &\text{ iff } \mathcal{A} \models g(x) < g(y) && \text{since } \mathcal{A} \models \text{SUBSET, INDISC}^\sim \\ &\text{ iff } \hat{\mathcal{A}} \models g(x) < g(y) && \text{since } \hat{\mathcal{A}} \subseteq \mathcal{A} \end{aligned}$$

e.

$$\begin{aligned}
\mathcal{Q}_f \models \text{Sum}(x, y, z) &\text{ iff } \mathcal{Q}_f \models (x' \cup y' = z \\
&\quad \wedge x \sim x' \wedge y \sim y' \\
&\quad \wedge x' \cap y' = \emptyset) && \text{since } \mathcal{Q}_f \models \text{DEF}^+ \\
&\text{ iff } \hat{\mathcal{A}} \models (g(x') \cup g(y') = g(z) \\
&\quad \wedge g(x) \sim g(x') \\
&\quad \wedge g(y) \sim g(y') \\
&\quad \wedge g(x') \cap g(y') = \emptyset) && \text{by (b) and (c)} \\
&\text{ iff } \mathcal{A} \models (g(x') \cup g(y') = g(z) \\
&\quad \wedge g(x) \sim g(x') \\
&\quad \wedge g(y) \sim g(y') \\
&\quad \wedge g(x') \cap g(y') = \emptyset) \\
&\text{ iff } \mathcal{A} \models \text{Sum}(g(x), g(y), g(z)) && \text{since } \mathcal{A} \models \text{DEF}^+ \\
&\text{ iff } \hat{\mathcal{A}} \models \text{Sum}(g(x), g(y), g(z)) && \text{since } \hat{\mathcal{A}} \subseteq \mathcal{A}
\end{aligned}$$

□

So, each model of CAI_f has a submodel isomorphic to $\mathcal{Q}_f$, and we may conclude:

Corollary 5.2.27. *If f is total and congruous, then CAI_f has a prime model.*

5.3 Summary

We can now draw our final conclusions about CA, CS, and their completions.

Theorem 5.3.1. *If f is a total, congruous remainder function, then CAI_f is consistent and complete.*

Proof. CAI_f is consistent because f is congruous, by 3.3.12c. Since CAI_f is model complete, by 5.1.6, and has a prime model, by 5.2.27, the prime model test (Appendix, Fact C.3) applies. So CAI_f is complete. □

Corollary 5.3.2. *If T is a completion of CAI, then $T \equiv \text{CAI}_f$, for some congruous f .*

Proof. For each $n > 0$, $T \vdash \text{MOD}_{n,i}$ for exactly one i , $0 \leq i < n$:

Since T is complete, $T \vdash \text{MOD}_{n,i}$ or $T \vdash \neg \text{MOD}_{n,i}$ for each such i . But if $T \vdash (\neg \text{MOD}_{n,0} \wedge \dots \wedge \neg \text{MOD}_{n,n-1})$, then T is inconsistent, since $T \vdash \text{CAI}$ and $\text{CAI} \vdash \text{DIV}_n$. Hence $T \vdash \text{MOD}_{n,i}$ for at least one i , $0 \leq i < n$.

But suppose $T \vdash \text{MOD}_{n,i}$ and $T \vdash \text{MOD}_{n,j}$ where $0 \leq i \neq j < n$. Again, T would be inconsistent, for $\text{CA} \vdash \text{MOD}_{n,i} \rightarrow \neg \text{MOD}_{n,j}$ (see Lemma 3.3.11).

So, let $f(n) = m$ iff $T \vdash \text{MOD}_{n,m}$. Then $T \vdash \text{CAI}_f$ and, since CAI_f is complete, $\text{CAI}_f \vdash T$. □

Corollary 5.3.3.

- a. Every completion of CA is consistent with CS.
- b. $CA \equiv CS$

Proof.

- a. Follows from 5.3.2 and 3.3.13.
- b. Follows from (a) and 3.3.2b, given that $CS \vdash CA$.

□

Theorem 5.3.4.

- a. For $n > 0$, $CS; EXACTLY_n$ is decidable.
- b. CS is decidable.

Proof.

- a. $CS; EXACTLY_n \vdash \phi$ iff $\mathcal{A}_n \models \phi$. But $\mathcal{A}_n$ is a finite model.
- b. To determine whether $CS \vdash \phi$, alternate between generating theorems of CA and testing whether $\mathcal{A}_n \models \neg\phi$.

□

Corollary 5.3.5.

- a. CSI has 2^ω completions.
- b. For total f , CSI_f is decidable iff f is decidable.

Proof.

- a. There are 2^ω remainder functions whose domain is the set of prime numbers. Each such function is congruous, so each corresponds to a consistent extension of CSI. By Lindenbaum's lemma, each of these extensions has a consistent and complete extension.
- b. ($\Rightarrow$) If f is decidable, then CAI_f is recursively enumerable. But CAI_f is complete, so it is decidable.
 ($\Leftarrow$) To calculate $f(n)$, determine which $MOD_{n,m}$ sentence is in CSI_f .

□

Theorem 5.3.6. *There is no sentence, ϕ , such that $T = CA; \phi$ is consistent and T has only infinite models.*

Proof. If ϕ is true only in infinite models of CA, then $\neg\phi$ is true in all finite models of CA, so $\neg\phi \in \text{CS}$. But $CA \equiv \text{CS}$, so $CA; \phi$ is inconsistent. $\square$

Theorem 5.3.7. *CS is not finitely axiomatizable.*

Proof. Suppose $\text{CS} \vdash \phi$. So $CA \vdash \phi$ and, by compactness, $(\text{BASIC} \cup T) \vdash \phi$ for some finite set of ADIV_n principles, $T = \{\text{ADIV}_n \mid n \in J\}$. Let $K = \{n \mid \text{every prime factor of } n \text{ is a member of } J\}$.

Let $\mathcal{A}$ be a model with domain $\bigcup_{k \in K} \mathbf{Q}_k$, in which size relations are determined in accordance with the size function, θ , defined in 3.2.26. We claim the following without proof:

- (1) $\mathcal{A} \models \text{BASIC}$
- (2) $\mathcal{A} \models \text{ADIV}_j$ for all $j \in J$
- (3) $\mathcal{A} \not\models \text{ADIV}_k$

By (1) and (2), $\mathcal{A} \models (\text{BASIC} \cup T)$, so $\mathcal{A} \models \phi$. But by (3), $\mathcal{A} \not\models \text{CS}$. Hence $\phi \notin \text{CS}$. $\square$

6

Sets of Natural Numbers

CS has standard finite models since it consists of sentences true in all such models. It has infinite models $\mathcal{Q}_n$ and $\mathcal{Q}_f$. In this chapter we will show that CS has infinite standard models over $\mathcal{P}(\mathbb{N})$.

An ordering of $\mathcal{P}(\mathbb{N})$ that satisfies CS will not necessarily appear reasonable. For example, some such orderings say that there are fewer even numbers than prime numbers (see below 6.2.13). To rule out such anomalies, we introduce a principle, OUTPACING, in section 1. OUTPACING mentions the natural ordering of $\mathbb{N}$ and applies only to subsets of $\mathbb{N}$. Section 2 establishes that OUTPACING can be satisfied jointly with any consistent extension of CS in a model whose domain is $\mathcal{P}(\mathbb{N})$. So CS ; OUTPACING does not fix the size relations over $\mathcal{P}(\mathbb{N})$.

6.1 The outpacing principle

Throughout this chapter, x and y will range over $\mathcal{P}(\mathbb{N})$.

Definition 6.1.1. x *outpaces* y just in case the restriction of x to any sufficiently large initial segment of $\mathbb{N}$ is larger than the corresponding restriction of y , that is, iff:

$$\exists n \forall m (m > n \rightarrow |x^{[m]}| > |y^{[m]}|)$$

Notice that the size comparison between the two restricted sets will always agree with the comparison of their normal cardinalities since all initial segments of $\mathbb{N}$ are finite.

We employ this notion to state a *sufficient* condition for one set of natural numbers to be larger than another:

OUTPACING. If x outpaces y , then $x \not\prec y$.

The general motivation behind this principle should be familiar. We extrapolate from well understood finite cases to puzzling infinite cases. But we should also

emphasize, again, that this extrapolation cannot be done in any straightforward, mechanical way without risking contradiction. We cannot, for example, strengthen the conditional to a biconditional, thus:

$$(1) \quad x \dot{\prec} y \text{ iff } x \text{ outpaces } y.$$

This revised principle conflicts with CS, for *outpacing* is not a quasi-linear ordering. For example, neither $[2\mathbf{n}]$ nor $[2\mathbf{n} + 1]$ outpaces the other since each initial segment $\{0, \dots, 2n + 1\}$ of $\mathbb{N}$ contains n evens and n odds. But the two are discernible under *outpacing*, since $[2\mathbf{n}]$ outpaces $[2\mathbf{n} + 2]$ while $[2\mathbf{n} + 1]$ does not.

There is another point that underlines the need for care in extrapolating from finite cases to infinite cases: we cannot just use (2):

$$(2) \quad \text{If, given any finite subset } z \text{ of } \mathbb{N}, x \text{ restricted to } z \text{ is larger than } y \text{ restricted to } z, \text{ then } x > y.$$

Though (2) is true, its antecedent is only satisfied when $y \subset x$.

So, there are many statements that assert of infinite cases what is true of finite cases. Some of these conflict with one another. Others are too weak to be helpful. It is doubtful whether there is any mechanical way to decide which of these statements are true. The best we can do is propose plausible theories, determine whether they are consistent, and see how far they go.

Definition 6.1.0. *$\mathcal{A}$ is an **outpacing model** iff*

$$\begin{aligned} \dot{\mathcal{A}} &= \mathcal{P}(\mathbb{N}) \\ \mathcal{A} &\models \text{BASIC, and} \\ \mathcal{A} &\models \text{OUTPACING} \end{aligned}$$

There is a slight difficulty in saying that an interpretation of $L_{C<}$ satisfies *OUTPACING*. Since *OUTPACING* involves the $\dot{\prec}$ relation over $\mathbb{N}$, it cannot be expressed in $L_{C<}$. We shall finesse this problem by regarding *OUTPACING* as the (very large) set of sentences

$$\{ \mathbf{b} < \mathbf{a} \mid a \text{ outpaces } b \}$$

Fact 6.1.1. *Every outpacing model satisfies the following*

- a. $[2\mathbf{n}] > [3\mathbf{n}]$
- b. $[3\mathbf{n}] > [4\mathbf{n}] > [5\mathbf{n}] > \dots$
- c. If $k > 0$, then $[k\mathbf{n}] > [\mathbf{n}^2]$

Proof.

- a. $[2\mathbf{n}]$ has at least $(k-1)/2$ members less than or equal to k for any given k .
 $[3\mathbf{n}]$ has at most $k/3 + 1$ such members. If $k > 4$, then $(k-1)/2 > k/3 + 1$

- b. Similar to (1).
- c. If $m = k^2$, both $[k\mathbf{n}]$ and $[\mathbf{n}^2]$ have exactly k members less than m . For $m > k(k+1)$, $\mathbb{N}^{[m]}$ will have more members in $[k\mathbf{n}]$ than in $[\mathbf{n}^2]$.

□

Fact 6.1.2. *Every outpacing model satisfies the following*

- a. $[2\mathbf{n}] \geq [2\mathbf{n} + 1]$
- b. $[2\mathbf{n} + 1] \geq [2\mathbf{n} + 2]$
- c. If $[2\mathbf{n}] > [2\mathbf{n} + 1]$, then $[2\mathbf{n} + 1] \sim [2\mathbf{n} + 2]$.

Proof.

- a. By TRICH, it is sufficient to show that $\neg[2\mathbf{n}] < [2\mathbf{n} + 1]$. If it were, then, by $\text{REP}^<$, there is a y such that $y \sim [2\mathbf{n}]$ and $y \subset [2\mathbf{n} + 1]$. But any proper subset of $[2\mathbf{n} + 1]$ is outpaced by, and hence smaller than $[2\mathbf{n}]$. Let k be the least odd number not in y . Then $[2\mathbf{n}]$ leads y at $k+1$ and y never catches up. So there is no y such that $[2\mathbf{n}] \sim y \subset [2\mathbf{n} + 1]$.
- b. Similar to that of (1).
- c. Note that $[2\mathbf{n}] = [2\mathbf{n} + 2] \cup \{0\}$ and that $\text{BASIC} \models (*)$.

$$(y \subset x \wedge z \leq y \wedge \text{Atom}(z') \wedge z \subset x \wedge x = z \cup z') \rightarrow y \sim z$$

Let $x = [2\mathbf{n}]$, $y = [2\mathbf{n} + 1]$, $z = [2\mathbf{n} + 2]$ and apply $(*)$.

□

The two alternatives left open in 6.1.2 correspond to the possibilities that $\mathbb{N}$ may be odd or even: if $[2\mathbf{n}] \sim [2\mathbf{n} + 1]$, then $\mathbb{N}$ is even, if $[2\mathbf{n} + 1] \sim [2\mathbf{n} + 2]$, then $\mathbb{N}$ is odd. In section 6.2, we show that both of these possibilities can be realized in standard models over $\mathcal{P}(\mathbb{N})$. Here, we generalize 6.1.2 to similar cases, including other congruence classes.

Definition 6.1.3. $\langle x, y \rangle$ is an **alternating pair** iff x and y are infinite and for all $i > 0$, $x_i < y_i < x_{i+1}$.

Fact 6.1.4. *If $\langle x, y \rangle$ is an alternating pair, then in any outpacing model:*

$$x \sim y \vee x > y \sim (x - x_1)$$

Proof. The argument for 6.1.2 applies here since the only facts about $[2\mathbf{n}]$ and $[2\mathbf{n} + 1]$ used hold by virtue of these sets forming an alternating pair. □

Theorem 6.1.5. *For a given $k > 0$, let $A_i = [k\mathbf{n} + i]$ for each $i < k$. Then:*

a. If $0 \leq i < j < k$, then $\langle A_i, A_j \rangle$ is an alternating pair.

b. There is a p , $0 < p \leq k$ such that

(i) If $i < j < p$, then $A_i \sim A_j$.

(ii) If $p \neq k$, then $A_0 \sim A_p \cup \{0\}$.

(iii) If $p \leq i < k$, then $A_i \sim A_p$.

(See example below).

Proof.

a. $A_i(n) = kn + i, A_j(n) = kn + j, A_i(n+1) = kn + i + k$ and $i < j < k + i$.

b. If $A_0 > A_i$ for some i , let p be the least such i ; otherwise, let $p = k$.

(i) For $0 \leq i < p$, $\langle A_0, A_i \rangle$ is an alternating pair. So either $A_0 > A_i$ or $A_0 \sim A_i$ by 6.1.4. But $A_0 \leq A_i$ by the selection of p , so $A_0 \sim A_i$. (i) follows by TRANS~.

(ii) Immediate from 6.1.4 since $\langle A_0, A_p \rangle$ is an alternating pair and $A_0 > A_p$.

(iii) $A_0 > A_p \geq A_i$ if $i \geq p$. So $A_0 > A_i$. Hence $A_i \sim A_0 - \{0\} \sim A_p$. So $A_i \sim A_p$.

□

Example. Let $k = 4$, so $A_i = [4\mathbf{n} + i]$ for $i = 0, 1, 2, 3$. Then one of the following situations obtains:

$$A_0 \sim A_1 \sim A_2 \sim A_3 > [4\mathbf{n} + 4]$$

$$A_0 > A_1 \sim A_2 \sim A_3 \sim [4\mathbf{n} + 4]$$

$$A_0 \sim A_1 > A_2 \sim A_3 \sim [4\mathbf{n} + 4]$$

$$A_0 \sim A_1 \sim A_2 > A_3 \sim [4\mathbf{n} + 4]$$

6.2 Models of CS and Outpacing

In this section, we construct models of CS over $\mathcal{P}(\mathbb{N})$ that satisfy OUTPACING.

Outpacing models will be constructed out of finite models of CS by a technique which is very much like the ultraproduct construction common in model theory, though the application here demands some important differences.

Definition 6.2.1.

a. L_N , **the language of subsets of $\mathbb{N}$** , is the first order language which results from adding to $L_{C<}$, as individual constants, a name for each subset of $\mathbb{N}$.

b. $\mathcal{A}_n$ is the standard finite interpretation of L_N over $\mathcal{P}(\mathbb{N}^{[n]})$ in which:

$$\mathcal{A}_n(\mathbf{a}) = a \cap \mathbb{N}^{[n]} = a^{[n]} \text{ for each } a \subseteq \mathbb{N}.$$

Definition 6.2.2. (cf. [Monk, Def. 18.15, p318]) If X is a set and $F \subseteq \mathcal{P}(X)$, then

- a. F has the **finite intersection property** iff the intersection of any finite subset of F is non-empty.
- b. F is a **filter** over X iff
 - (i) $F \neq \emptyset$
 - (ii) If $a \in F$ and $a \subseteq b$, then $b \in F$, and
 - (iii) If $a \in F$ and $b \in F$, then $a \cap b \in F$
- c. F is an **ultrafilter** over X iff
 - (i) F is a filter over X
 - (ii) $X \in F$, and
 - (iii) if $Y \subseteq X$, then either $Y \in F$ or $(X - Y) \in F$.
- d. An ultrafilter, F , over X is **principal** iff there is some $x \in X$ such that $F = \{a \subseteq X \mid x \in a\}$

Fact 6.2.3.

- a. A non-principal ultrafilter contains no finite sets. [Bell, Ch.6, lemma 1.3, p.108]
- b. A non-principal ultrafilter over X contains all cofinite subsets of X .
- c. If $F \subseteq \mathcal{P}(X)$ and F has the finite intersection property, then there is an ultrafilter over X which includes F . [Monk, Prop. 18.18, p.319]
- d. If $Y \subseteq X$ and Y is infinite, then there is a non-principal ultrafilter over X which contains Y .

Definition 6.2.4. If F is an ultrafilter over $\mathbb{N}$, then $\mathcal{A}_F$ is the interpretation of $L_{C<}$ in which

- a. $\dot{\mathcal{A}}_{\mathcal{F}} = \mathcal{P}(\mathbb{N})$,
- b. $\mathcal{A}_F \models \mathbf{a} < \mathbf{b}$ iff $\{k \mid a^{[k]} < b^{[k]}\} \in F$, and similarly for other predicates.
- c. Boolean symbols receive the usual interpretation.

Our main result is that if F is non-principal, then $\mathcal{A}_F$ is an outpacing model.

Theorem 6.2.5. *If F is a non-principal ultrafilter over $\mathbb{N}$, then*

a. *If τ is a term of L_N , then*

$$\mathcal{A}_k(\tau) = \mathcal{A}_F(\tau) \cap A_k = \mathcal{A}_F(\tau)^{[k]}$$

b. *If ϕ is a quantifier free formula of L_N , then*

$$\mathcal{A}_F \models \phi \text{ iff } \{k \mid \mathcal{A}_k \models \phi\} \in F$$

c. *If ϕ is a universal formula of L_N and $\mathcal{A}_k \models \phi$ for every k , then $\mathcal{A}_F \models \phi$.*

d. $\mathcal{A}_F \models REP^<$.

e. $\mathcal{A}_F \models BASIC$.

f. $\mathcal{A}_F \models ADIV_n$, for every n .

Proof.

a. By induction on the structure of τ :

(i) If τ is a constant, $\tau = a$ for some $a \in \mathbb{N}$. So $\mathcal{A}_k(\tau) = a^{[n]}$ by 6.2.1b.

(ii) If $\tau = \ulcorner \tau_1 \cup \tau_2 \urcorner$,

$$\begin{aligned} \mathcal{A}_k(\tau) &= \mathcal{A}_k(\tau_1) \cup \mathcal{A}_k(\tau_2) \\ &= (\mathcal{A}_F(\tau_1) \cap A_k) \cup (\mathcal{A}_F(\tau_2) \cap A_k) \\ &= (\mathcal{A}_F(\tau_1) \cup \mathcal{A}_F(\tau_2)) \cap A_k \\ &= \mathcal{A}_F(\tau_1 \cup \tau_2) \cap A_k \end{aligned}$$

The proofs for intersections and relative complements are similar.

b. By induction on the structure of ϕ :

(i) If $\phi = \ulcorner a \subset b \urcorner$, then $\mathcal{A}_F \models \phi$ iff $a \subset b$

If $a \subset b$, then there is a $k \in b$ but not $\in a$. So if $n > k$, $a^{[n]} \subset b^{[n]}$.

Hence, $\{n \mid \mathcal{A}_F \models \phi\}$ is cofinite and, by 4.2.3b, in F .

Conversely, if $\{n \mid a^{[n]} \subset b^{[n]}\} \in F$, then it is infinite. So, there cannot be a k in a but not in b ; otherwise $a^{[n]}$ would not be included in $b^{[n]}$ for any n greater than k . So $a \subseteq b$. But, clearly $a \neq b$, so $a \subset b$.

(ii) If $\phi = \ulcorner a = b \urcorner$, then

$$\begin{aligned} \mathcal{A}_F \models \phi &\text{ iff } a = b \\ &\text{ iff } a^{[n]} = b^{[n]} \text{ for all } n. \\ &\text{ iff } \{n \mid \mathcal{A}_n \models a = b\} = \mathbb{N} \\ &\text{ iff } \{n \mid \mathcal{A}_n \models a = b\} \in F \end{aligned}$$

since if $\mathcal{A}_n \not\models a = b$ and $k > n$, then $\mathcal{A}_k \not\models a = b$.

(iii) $\mathcal{A}_F \models a < b$ iff $\{k \mid \mathcal{A}_k \models (a < b)\} \in F$. Immediate from 6.2.4b.

(iv) If ϕ is non-atomic, then, since F is an ultrafilter:

$$\begin{aligned} \mathcal{A}_F \models \phi_1 \wedge \phi_2 &\text{ iff } \mathcal{A}_F \models \phi_1 \text{ and } \mathcal{A}_F \models \phi_2 \\ &\text{ iff } \{k \mid \mathcal{A}_k \models \phi_1\} \in F \text{ and } \{k \mid \mathcal{A}_k \models \phi_2\} \in F \\ &\text{ iff } \{k \mid \mathcal{A}_k \models \phi_1 \wedge \phi_2\} \in F \end{aligned}$$

$$\begin{aligned} \mathcal{A}_F \models \neg\phi &\text{ iff } \mathcal{A}_F \not\models \phi \\ &\text{ iff } \{k \mid \mathcal{A}_k \models \phi\} \notin F \\ &\text{ iff } \{k \mid \mathcal{A}_k \models \neg\phi\} \in F \end{aligned}$$

c.

Suppose $\mathcal{A}_k \models \forall x\phi(x)$, for all k
 then $\mathcal{A}_k \models \phi(a)$, for all a , for all k
 So $\mathcal{A}_k \models \phi(a)$, for all k , for all a
 So $\mathcal{A}_F \models \phi(a)$, for all a by (b)
 So $\mathcal{A}_F \models \forall x\phi(x)$

d. Suppose $\mathcal{A}_F \models (a < b)$. Construct a' , a subset of b , for which $\mathcal{A}_F \models (a \sim a')$ as follows:

Let $K = \{k \mid a^{[k]} < b^{[k]}\}$ so, $K \in F$.
 $= \{k_1, \dots, k_i, \dots\}$ where the k_i 's are in strictly increasing order.

Let $a_0 = \emptyset$

$$a_{i+1} = a_i \cup \left\{ \text{the } n \text{ greatest members of } b^{[k_{i+1}]} \text{ not in } a_i \right\},$$

$$\text{where } n = |a^{[k_{i+1}]}| - |a_i|$$

$$a' = \bigcup a_i$$

Then $a' \subset b$ since each a_i draws its new members from b .

Claim: If $k \in K$, then $a'^{[k]} \sim a^{[k]}$.

Hence: $\mathcal{A}_F \models a' \sim a$ since they are the same size over some set which contains K and is, thus, in F .

e. Immediate from (c) and (d): BASIC is equivalent to a set of universal sentences, together with ATOM and $\text{REP}^<$. ($\mathcal{A}_F \models \text{ATOM}$ because it contains all singletons of natural numbers.)

f. For $0 < i \leq n$, x an infinite subset of $\mathbb{N}$, let $x_i = \{x_{kn+i-1} \mid k \in \mathbb{N}\}$.

The n sets, x_i , partition x . Furthermore, these form an alternating n -tuple, in the manner of the congruence classes modulo n (see Theorem 6.1.5). As in 6.1.5, these sets are approximately equal in size and $\mathcal{A}_F \models \text{ADIV}_n(x)$.

□

Corollary 6.2.6. *If F is a non-principal ultrafilter over $\mathbb{N}$, then*

- a. $\mathcal{A}_F \models CAI$, and
- b. $\mathcal{A}_F \models CSI$

Proof.

- a. Immediate from 6.2.5e and 6.2.5f.
- b. Immediate from (a) and 5.3.3b.

□

The proof of 6.2.6 is modeled on the usual ultraproduct construction, but is not quite the same. In the usual construction(see, for example, [Bell, pp. 87-92]), a model is built by first taking the product of all factors (in this case, the $\mathcal{A}_k$), which results in a domain whose elements are functions from the index set ($\mathbb{N}$, here) to elements of the factors. These functions are then gathered into equivalence classes (by virtue of agreeing almost everywhere, i.e. on some member of the filter) and the reduced ultraproduct is defined by interpreting the language over these equivalence classes. The model so constructed, which we'll call $\prod \mathcal{A}_i/F$, has the handy property that it satisfies any formula which is satisfied by almost all factors, and certainly any formula which is satisfied in all the factors. This is handy because, given that each of the $\mathcal{A}_k$ satisfies CS, we can immediately conclude that $\prod \mathcal{A}_i/F$ satisfies CS.

Unfortunately, $\prod \mathcal{A}_i/F$ is not the model we wanted: its elements are not subsets of $\mathbb{N}$, but equivalence classes of functions from $\mathbb{N}$ to finite subsets of $\mathbb{N}$. There is, indeed, a natural mapping from subsets of $\mathbb{N}$ to such elements, and this mapping would have allowed us to identify a model over $\mathcal{P}(\mathbb{N})$ as a submodel of $\prod \mathcal{A}_i/F$; but only a submodel. So, had we constructed the reduced ultraproduct, we would have then been able to infer that the part of that model which held our interest satisfied all universal formulas of CS; we still would have had to resort to special means to show that the non-universal formulas were likewise satisfied.

Fortunately, these special means were available; the only non-universal axioms of CAI could be verified in the constructed model more or less directly, and the completeness proof of the last chapter allowed us to infer that all formulas true in all of the factors are true in the model $\mathcal{A}_F$, after all.

For a given F , there will be many cases where $\mathcal{A}_F \models \mathbf{a} < \mathbf{b}$ even though b does not outpace a . This will happen whenever $\{k \mid a^{[k]} < b^{[k]}\} \in F$ but is not cofinite. Consider a familiar example: Let $a = [2\mathbf{n} + 1]$ and $b = [2\mathbf{n}]$. Then $a^{[k]} < b^{[k]}$ iff $k \in [2\mathbf{n}]$, for if we count the even numbers and the odd numbers up to some even number, there will always be one more even and if we count up to some odd number, there will always be the same number of evens and odds.

$[2\mathbf{n}]$ is neither finite nor cofinite, so it may or may not be in F . If $[2\mathbf{n}] \in F$, $\mathcal{A}_F \models [2\mathbf{n} + 1] < [2\mathbf{n}]$. Otherwise, $[2\mathbf{n} + 1] \in F$, so $\mathcal{A}_F \models [2\mathbf{n} + 1] \sim [2\mathbf{n}]$.

The construction of a model $\mathcal{A}_F$ from any non-principal ultrafilter, F , suggests that there are many outpacing models unless different ultrafilters can yield the same model. We will first show that this qualification is not needed.

Theorem 6.2.7. *If F_1 and F_2 are distinct non-principal ultrafilters over $\mathbb{N}$, then $\mathcal{A}_{F_1} \neq \mathcal{A}_{F_2}$.*

Proof. (See below.) □

To show this, we will show that the presence of a set in an ultrafilter makes a direct, personalized contribution to the model $\mathcal{A}_F$. Putting this in another way, there is a set of decisions that must be made in constructing an outpacing model; each decision may go either way, though the decisions are not independent of each other. Furthermore, each decision is made for a model $\mathcal{A}_F$ by the presence or absence of a particular set in F .

Definition 6.2.8. *If $x \subseteq \mathbb{N}$, then $x^+ = \{i + 1 \mid i \in x\}$.*

A pair of sets, $\langle x, x^+ \rangle$, can sometimes be an alternating pair, but this is not always the case.

Fact 6.2.9. *$\langle x, x^+ \rangle$ is an alternating pair iff x is infinite and there is no $n \in x$ such that $(n + 1) \in x$. That is, no two consecutive numbers are in x .*

Nevertheless, pairs $\langle x, x^+ \rangle$ are like alternating pairs in the following way:

Lemma 6.2.10. *If x is infinite, $x_1 = (x - x^+)$, $x_2 = (x^+ - x)$, F is a non-principal ultrafilter, and $\mathcal{A} = \mathcal{A}_F$, then*

- a. $\langle x_1, x_2 \rangle$ is an alternating pair.
- b. $\mathcal{A} \models x \sim x^+$ or $\mathcal{A} \models x > x^+ \sim x - \{x_1\}$
- c. $\mathcal{A} \models x > x^+$ iff $x \in F$.

Proof.

- a. Let a **run** of x be a maximal consecutive subset of x . (So $[2\mathbf{n}]$ has only 1-membered runs, while $\mathbb{N} - [10\mathbf{n} + 1]$ has only 9-membered runs.) So, $x_1(n)$ is the first element in the n -th run of x and $x_2(n)$ is the first element after the n -th run of x .
- b. We know from (a) that $x_2 \sim x_1$ or $x_2 \sim x_1 - x_1(1)$. But the disjoint union of $x \cap x^+$ with x_1 or x_2 , respectively, yields x or x^+ . So (b) follows from $\text{DISJ}^\cup$.
- c. We need only prove (*):

$$(*) \quad |x^{[n]}| > |x^{+[n]}| \text{ iff } n \in x$$

Informally: $x^{[n]}$ first becomes greater than $x^{+[n]}$ for $n = x(1)$, since $x(1) \notin x^+$ because $\neg(x_1 - x_1(1) \subset x)$. Throughout the first run of x , x maintains its lead, losing it only at the least n for which $n \notin x$ (since $(n-1) \in x$, so $n \in x^+$). This pattern repeats during successive runs of x .

□

We can now prove our main result, Theorem 6.2.7:

Proof. Without loss of generality, we can suppose there is a set, x , such that $x \in F_1$ and $x \notin F_2$. By 6.2.10c, $\mathcal{A}_{F_1} \models x > x^+$ and $\mathcal{A}_{F_2} \models x \sim x^+$ □

Theorem 6.2.7 allows us to improve upon some previous results. For example, we can show that either of the alternatives in 6.2.10b can be obtained for any alternating pair.

Theorem 6.2.11.

- a. If neither x nor y outpaces the other, then there is an ultrafilter F , such that $\mathcal{A}_F \models (x \sim y)$.
- b. If $\langle x, y \rangle$ is an alternating pair, then there is an ultrafilter F , such that $\mathcal{A}_F \models (x > y)$.

Proof.

- a. Let $J = \{k \mid |x^{[k]}| = |y^{[k]}|\}$. Since neither x nor y outpaces the other, J is infinite. By 6.2.3d, let F be a non-principal ultrafilter which contains J . Then $\mathcal{A}_F \models x \sim y$.
- b. If $\langle x, y \rangle$ is an alternating pair, so is $\langle y, x - x(1) \rangle$. So, by (a) there is an F such that $\mathcal{A}_F \models y \sim (x - x(1))$. But then $\mathcal{A}_F \models x > y$.

□

Theorem 6.2.12. Every infinite completion of CS has an outpacing model.

Proof. Recall that every infinite completion of CS is equivalent to CSI_f for some total and congruous remainder function, f . (See 3.6.2.)

Given such an f , let $G_k = [k\mathbf{n} + f(k) + 1]$ for each $k > 0$. Then (*) holds for each k :

$$(*) \quad G_k = \{n \mid \mathcal{A}_n \models \text{MOD}_{k, f(k)}\}$$

Let $G = \{G_k \mid k > 0\}$.

The intersection of any finite subset, H , of G is infinite. If H is a finite subset of G , then $H = \{G_k \mid k \in J\}$, where J is some finite subset of $\mathbb{N}^+$. So $H = \{n + 1 \mid k \in J \rightarrow n \equiv f(k) \pmod{k}\}$. But f is congruous, so the restriction of f to the finite domain J has infinitely many solutions (see 3.3.8a).

Since the intersection of any finite subset of G is infinite, there is a non-principal ultrafilter F such that $G \subseteq F$. By (*), $\mathcal{A}_F \models \text{MOD}_{k, f(k)}$, so $\mathcal{A}_F \models \text{CSI}_f$. □

On the basis of 6.2.11 we noted that there are even and odd outpacing models; we can now extend that observation to moduli other than 2. More specifically, all of the possibilities listed in 6.1.5 for the relative sizes of the k -congruence classes are obtainable in outpacing models.

This section has explored the existence and variety of outpacing models. Three comments are in order before we turn to the common structure of outpacing models.

First, even if T is an infinite completion of CS, there is no unique outpacing model which satisfies T . This would be true only if fixing the congruence classes determined whether $x \sim x^+$ or $x > x^+$ for every $x \subseteq N$. That all such choices are not determined by a remainder theory can be seen intuitively, perhaps, by considering $x = [\mathbf{n}^2]$: any finite set of congruences has infinitely many solutions that are squares and infinitely many that are not; so whether $x \in F$ is an independent choice. Also, there are only 2^ω remainder functions while there are 2^{2^ω} non-principal ultrafilters over $\mathbb{N}$ [Bell, Ch. 6, Theorem 1.5], each yielding a different outpacing model.

Second, it is not clear whether every outpacing model can be obtained by the construction of 6.2.4. Lemma 6.2.10c may suggest that any outpacing model, $\mathcal{A}$, is $\mathcal{A}_F$ for $F_{\mathcal{A}} = \{x \mid \mathcal{A} \models x > x^+\}$, but it should not. To establish that $\mathcal{A} = \mathcal{A}_F$, both (1) and (2) are necessary.

- (1) If $\mathcal{A}$ is an outpacing model, then $F_{\mathcal{A}}$ is a non-principal ultrafilter.
- (2) If $F_{\mathcal{A}} = F_{\mathcal{B}}$, then $\mathcal{A} = \mathcal{B}$

I have not been able to prove (1) or (2). If (1) is false, then clearly $\mathcal{A} \neq \mathcal{A}_F$. But even if (1) is true, two outpacing models may agree about all pairs $\langle x, x^+ \rangle$ but disagree elsewhere. At most one of them is obtainable by our construction. So (1) and (2) are open problems.

Finally, though it may be extraneous to show that OUTPACING is independent, we will do so.

Theorem 6.2.13. *There are standard models of CS over $\mathcal{P}(\mathbb{N})$ which do not satisfy OUTPACING.*

Proof. Suppose that $\prec$ is a linear ordering under which $\mathbb{N}$ forms an ω -sequence. Then we could define $x \prec$ -outpaces y as follows:

$$\exists n \forall m [n \prec m \rightarrow |\{k \mid k \in x \wedge k \prec m\}| > |\{k \mid k \in x \wedge k \prec n\}|]$$

and define the principle:

$$\text{OUTPACING}_{\prec} . \text{ If } x \prec\text{-outpaces } y, \text{ then } x > y.$$

Modifying 6.2.4, we could produce standard models of CS over $\mathcal{P}(\mathbb{N})$ which satisfy $\text{OUTPACING}_{\prec}$ and these will not, in general, satisfy OUTPACING.

Suppose, for example, that $\triangleleft$ is the ordering:

$$p_1, q_1, \dots, p_k, q_k, \dots$$

where p is the set of prime numbers and q is its complement. In any model of $\text{OUTPACING}_{\triangleleft}$, p and q will be nearly the same size, as the evens and the odds are in normal outpacing models. But the evens are much smaller than q , so $p > [2\mathbf{n}]$ and OUTPACING is false in $\text{OUTPACING}_{\triangleleft}$ models. $\square$

6.3 Size and density

When number theorists talk about the sizes of sets of natural numbers, they do not content themselves with speaking of the (Cantorian) cardinalities of these sets. Since they often want to compare infinite subsets of $\mathbb{N}$, they need a more discriminating notion.

One notion they use is **asymptotic density**. The asymptotic density of a set, x , of natural numbers is the limit, if there is one, of $|x^{[n]}|$ as n grows. For example, the asymptotic density of $[2\mathbf{n}]$ is $1/2$. From now on we shall use the term ‘density’ for asymptotic density.

In this section, we compare the ordering of $\mathcal{P}(\mathbb{N})$ given by density to the orderings given by CS and OUTPACING .

Definition 6.3.1.

- a. $\hat{x}^i = \frac{|x^{[i]}|}{i}$, the fraction of numbers less than or equal to i that are members of x .
- b. If $x \subseteq y \neq \emptyset$, then $\rho(x, y)$, the **density** of x in y , is the limit, if it exists of

$$\lim_{i=y_1}^{\omega} \hat{x}^i / \hat{y}^i$$

That is,

$$\rho(x, y) = r \text{ iff } \forall (\delta > 0) \exists n (\forall i > n) (r - \delta < \frac{\hat{x}^i}{\hat{y}^i} < r + \delta))$$

- c. The **density** of x , $\rho(x)$, is the density of x in $\mathbb{N}$, if x has a density in $\mathbb{N}$.
- d. If $x \subseteq y \neq \emptyset$, then x **converges** in y , $\text{cvg}(x, y)$, iff x has a density in y ; otherwise, x **diverges** in y .
- e. x **converges**, $\text{cvg}(x)$, iff x converges in $\mathbb{N}$. x **diverges** iff x diverges in $\mathbb{N}$.

Fact 6.3.2.

- a. If x is finite, $\rho(x) = 0$.
- b. If x is cofinite, $\rho(x) = 1$.
- c.

$$\begin{aligned}\rho([2\mathbf{n}]) &= 1/2 \\ \rho([4\mathbf{n}], [2\mathbf{n}]) &= 1/2 \\ \rho([4\mathbf{n}]) &= 1/4\end{aligned}$$

- d. If $\text{cvg}(x, y)$ and $\text{cvg}(y, z)$, then $\text{cvg}(x, z)$ and $\rho(x, z) = \rho(x, y)\rho(y, z)$.

Fact 6.3.3.

- a. There are divergent sets.
- b. If $0 \leq r \leq 1$, there is a set with density r .
- c. If $0 \leq r \leq 1$ and y is infinite, there is a set with density r in y .

Proof.

- a. Let $x = \{i \mid \exists n(10^{2n} \leq i < 10^{2n+1})\}$. So x contains all numbers between 0 and 9, between 100 and 999, between 10,000 and 99,999, and so forth. If $n > 1$, then $\hat{x}^{10^{2n}} \leq .1$ and $\hat{x}^{10^{2n+1}} \geq .9$. So $\hat{x}^k$ cannot have a limit.
- b. Suppose r is given. Construct the set x as follows:

$$x_0 = \emptyset$$

$$x_{i+1} = \begin{cases} x_i, & \text{if } \hat{x}^i \geq r; \\ x_i; i+1, & \text{otherwise.} \end{cases}$$

$$x = \bigcup x_i$$

- c. Modify the construction for (b) in the obvious ways.

□

Theorem 6.3.4. Suppose that both x and y converge in z . Then, if $\rho(x, z) < \rho(y, z)$, y outpaces x .

Proof.

Let $b = (\rho(y, z) - \rho(x, z))/3$
 Let $n_1 =$ the least n such that for $i \leq n$, $-b < \hat{x}^i/\hat{z}^i - \rho(x, z) < b$
 and let $n_2 =$ the least n such that for $i \leq n$, $-b < \hat{y}^i/\hat{z}^i - \rho(y, z) < b$
 So for any $i > n_1 + n_2$,
 we have $\hat{x}^i/\hat{z}^i < \rho(x, z) + b$
 and $\hat{y}^i/\hat{z}^i > \rho(y, z) - b$
 But $\rho(x, z) + b < \rho(y, z) - b$,
 So $\hat{x}^i/\hat{z}^i < \hat{y}^i/\hat{z}^i$
 So $\hat{x}^i < \hat{y}^i$
 So $|x^{[i]}| < |y^{[i]}|$

□

We can use the relation between density and outpacing to draw conclusions about densities that have nothing to do with outpacing, as in Theorem 6.3.5.

Theorem 6.3.5. *If $\rho(x, z_1) < \rho(y, z_1)$, and both x and y converge in z_2 , then $\rho(x, z_2) \leq \rho(y, z_2)$*

Proof. Since $\rho(x, z_1) < \rho(y, z_1)$, y outpaces x . But if $\rho(x, z_2) > \rho(y, z_2)$, then x outpaces y . So, $\rho(x, z_2) \leq \rho(y, z_2)$. □

We cannot strengthen the consequent of 6.3.5 to say that $\rho(x, z_2) < \rho(y, z_2)$: Let z_1 be the set of primes, let x contain every third member of z_1 , and let y be $z_1 - x$. Then $\rho(x, z_1) = 1/3$ and $\rho(y, z_1) = 2/3$, but $\rho(x, \mathbb{N}) = \rho(y, \mathbb{N}) = 0$.

Theorem 6.3.4 implies that in any outpacing model, sets with distinct densities will have distinct sizes. Even if two sets have the same density, they will differ in size if they have the same density in some common set. So, from Facts 6.3.b and (c), we can begin to appreciate how precise an ordering outpacing models provide:

Fact 6.3.6. *If $\mathcal{A}$ is an outpacing model, then*

- a. *there are uncountably many sizes of sets in $\mathcal{A}$, and*
- b. *if $0 \leq r \leq 1$, then even among sets with density r , there are uncountably many sizes in $\mathcal{A}$.*

Proof.

- a. Immediate from Fact 6.3.3b and Theorem 6.3.4.
- b. Let x be an infinite set with density r , let y be an infinite subset of x , where $\rho(y, x) = 0$, and let $z = x - y$. There are uncountably many subsets of y with distinct densities in y , though $\rho(y_1, x) = 0$, for any $y_1 \subset y$.

Suppose now that $y_1 \subset y$, $y_2 \subset y$, and $\rho(y_1) < \rho(y_2)$. Then $\mathcal{A} \models y_1 < y_2$, by 6.3.4. so $\mathcal{A} \models z \cup y_1 < z \cup y_2$, by $\text{DISJ}^\cup$.

But $\rho(z \cup y_1) = \rho(z \cup y_2) = \rho(x)$, since z was obtained by removing from x a set with density 0 relative to x .

□

6.3.1 The Convexity Problem

It's tempting to infer from these results that the extremely fine ordering of sets by size (or, rather, any such ordering which is realized in an outpacing model) is both a refinement and a completion of the ordering suggested by density: a refinement because it preserves all differences in size which are captured by the notion of density, a completion because all sets are located in a single, linear ordering of sizes.

But the situation is really not so clear. It is evident that the size ordering over $\mathcal{P}(\mathbb{N})$ in any outpacing is a refinement of the ordering by cardinality: if x has a smaller cardinal number than y , then x is smaller than y , though two sets with the same cardinal number may have different sizes. We can regard the cardinality of a set in $\mathcal{P}(\mathbb{N})$ as determined by its size, though different sizes may yield the same cardinal number. We shall express this fact by saying that, at least when we focus on $\mathcal{P}(\mathbb{N})$, cardinality is a function of size. (It is not at all clear that this is true in any power set.)

Now, we want to know whether the density of a set is a function of its size. It turns out that a negative answer is compatible with our theory (CS and OUTPACING), while an affirmative answer may or may not be consistent. First, we will give a more precise formulation of this problem; second, we will show that a negative answer is consistent; finally, we'll discuss the consistency of an affirmative answer. In passing, we'll explain why this is called *the convexity problem*.

Consider (1) and (1a):

$$(1) \quad \text{If } x \sim y \text{ and } \rho(x) = r, \text{ then } \rho(y) = r.$$

$$(1a) \quad \text{If } x \sim y \text{ and } \rho(x) = r \text{ and } \text{cvg}(y), \text{ then } \rho(y) = r.$$

(1a) is an immediate consequence of Theorem 6.3.4. For if y converges, there is some $r_2 = \rho(y)$; if $r < r_2$, then $x < y$ and if $r > r_2$, then $x > y$. But $x \sim y$, so $r_2 = r$.

So, the questionable part of (1) can be expressed as (2):

$$(2) \quad \text{If } x \sim y \text{ and } x \text{ converges, then } y \text{ converges.}$$

Theorem 6.3.4 insures that (1) just in case (2). Recalling that sets may have the same density even though they differ in size, we may consider two additional formulations:

$$(3) \quad \text{If } \rho(x) = \rho(y) \text{ and } x < z < y, \text{ then } \rho(z) = \rho(x)$$

(4) If $\rho(x) = \rho(y)$ and $x < z < y$, then z converges.

(3) and (4) are equivalent for the same reasons that (1) and (2) are equivalent.

Fact 6.3.7. *An outpacing model satisfies (1) iff it satisfies (3).*

Proof.

$\Rightarrow$ Suppose that $x < y < z$ and $\rho(x) = \rho(z)$.

By $\text{REP}^<$, there are two sets, x' and y' , for which

$$x' \subset y' \subset z, x \sim x', y \sim y'.$$

Since $x \sim x'$

and $\rho(x) = \rho(z)$

then $\rho(x') = \rho(z)$

by (1)

So $\rho(z - x') = 0$

and $\rho(z - y') = 0$

But $y' = z - (z - y')$

So $\rho(y') = \rho(z)$

So $\rho(y) = \rho(z)$

by (1)

$\Leftarrow$ Suppose that $x \sim y$ and $\rho(x) = r$.

If $x = \emptyset$ or $x = \mathbb{N}$, then $y = x$, so $\rho(y) = \rho(x)$

To apply (3), we need to find two sets, x_1 and x_2 such that

$$\rho(x_1) = \rho(x_2) = r = \rho(x)$$

and

$$x_1 < y < x_2$$

Assuming that $x \neq \emptyset$ and $x \neq \mathbb{N}$, let

$$x_1 = x - a, \text{ for some } a \in x$$

and

$$x_2 = x \cup b, \text{ for some } b \notin x.$$

Then $x_1 < x < x_2$, by SUBSET, and $x_1 < y < x_2$, by INDISC $^\sim$.

Since adding or removing a single element has no effect on the density of a set,

$$\rho(x_1) = \rho(x) = \rho(x_2)$$

So, by (3), $\rho(y) = r$.

□

The question at hand is called the ‘convexity problem’ because of the formulation in (3). In geometry, a figure is convex if, given any two points in the figure, any point between them (ie on the line segment from one to the other) is also in the figure. Applying this notion in the obvious way to the size ordering, (3) says that the class of sets having a given density is convex. By theorems 6.3.4 and 6.3.7, (1), (2) and (4) say the same thing.

We regret that the only thing we know about (3) is that it may be false:

Theorem 6.3.8. *The negation of (1) is satisfied in some outpacing model.*

Proof. Let x be a set with density r and let y be a divergent set which neither outpaces nor is outpaced by x . Let K be

$$\left\{ n \mid x^{[n]} = y^{[n]} \right\}$$

K is infinite, so K is a member of some non-principal ultrafilter, F . $\mathcal{A}_F \models x \sim y$, so (1) is false in $\mathcal{A}_F$. □

Open Problem: Is (1) consistent with CS and OUTPACING?

Appendix

A Notation

A.1 Predicate Logic

The formal theories discussed in this thesis are theories with standard formalization, in the sense of [Tarski, p.5]. That is, they are formalized in first order predicate logic with identity and function symbols. The following notation is used for the predicate calculus:

$\wedge$	and
$\vee$	or
$\neg$	not
$\rightarrow$	if ... then
$\leftrightarrow$	if and only if
$=$	identical
$\neq$	not identical
$\exists x$	there is an x
$\forall x$	for all x

Conjunctions and disjunctions of sets of sentences are represented by:

$$\bigwedge_{\dots \phi \dots} \phi$$

and

$$\bigvee_{\dots \phi \dots} \phi$$

A first order language is determined by its non-logical constant symbols, in the usual way. These may be predicates, individual constants, or function symbols. In most cases, the ranks of the symbols will accord with their familiar uses and we do not stipulate them.

A **schematic function** is a function whose range is a set of formulae. DIV_n , for example, maps natural numbers into sentences (see 3.2.8). The arguments of such functions are indicated by subscripts.

A.2 Set Theory

For first order languages with boolean operations and predicates, we use the following notation:

$\mathbf{I}$	the universe
$\emptyset$	the empty set
$x \cup y$	the union of x and y
$x \cap y$	the intersection of x and y
$x - y$	the relative complement of y in x
$x \subseteq y$	x is contained in y
$x \subset y$	x is a proper subset of y

These symbols are also used for set theoretic relations, outside of first order languages.

In addition, we use the following:

$x \in y$	x is a member of y
$\mathcal{P}(x)$	the power set of x
$\{x \mid \phi(x)\}$	the set of x 's which are ϕ
$x; a$	the union of x and a
$x^{[n]}$	the members of x less than or equal to n
$\mathbb{N}$	the set of natural numbers: 0, 1, ...
$\mathbb{N}^+$	$\mathbb{N} - 0$
$\mathbb{Z}$	the set of integers
ω	the smallest infinite cardinal
2^ω	2 to the ω

A.3 Arithmetic

For arithmetic, we use the following:

$i + j$	i plus j
$i - j$	i minus j
ij	i times j
i^j	i to the j -th power
$i!$	i factorial
$i \mid j$	i divides j
$\text{gcd}(i, j)$	greatest common divisor of i and j
$n \equiv m \bmod j$	n is congruent to m modulo j

We use the following notation for sets of natural numbers:

$$[\dots \mathbf{n} \dots] = \{ k \mid \exists n(k = \dots n \dots) \}$$

For example:

$$[k\mathbf{n} + j] = \text{the set of numbers congruent to } j \text{ modulo } k$$

$$[\mathbf{n}^2] = \text{the set of squares of natural numbers}$$

B Model Theory

This section lists the model-theoretic notions and results assumed in the text and presents our notation for these notions.

An **interpretation**, $\mathcal{A}$, of a first order language, L , consists of a domain, $\dot{\mathcal{A}}$, and a function which assigns to each individual constant of L a member of $\dot{\mathcal{A}}$, to each n -place predicate of L , a set of n -tuples over $\dot{\mathcal{A}}$, and to each n -argument function symbol of L , an n -ary function defined over and yielding values in $\dot{\mathcal{A}}$.

We assume the notion of **satisfaction** as defined in [Chang, section 1.3] and use

$$\mathcal{A} \models \phi \text{ to mean that } \mathcal{A} \text{ satisfies } \phi .$$

Definition B.1. *Familiar notions about models.*

- a. $\mathcal{A}$ is a **submodel** of $\mathcal{B}$ ($\mathcal{A} \subseteq \mathcal{B}$).
- b. $\mathcal{B}$ is an **extension** of $\mathcal{A}$ ($\mathcal{A} \subseteq \mathcal{B}$).
- c. The **submodel** of $\mathcal{B}$ generated by X , where X is a subset of $\dot{\mathcal{B}}$.
- d. $\mathcal{A}$ is **isomorphic** to $\mathcal{B}$ ($\mathcal{A} \simeq \mathcal{B}$).
- e. f is an **isomorphic embedding** of $\mathcal{A}$ into $\mathcal{B}$.
- f. $\mathcal{B}$ is an **elementary extension** of $\mathcal{A}$.
- g. $\{\mathcal{A}_i\}$ is a **chain** of models.
- h. $\mathcal{A} = \bigcup \{\mathcal{A}_i\}$; the **union of a chain** of models.

Definition B.2. *Notions about theories:*

- a. A **theory** is a set of first-order sentences.
- b. The **language of a theory**, L_T .
- c. T **proves** ϕ ($T \vdash \phi$);
 T **proves** T_2 ($T \vdash T_2$).

- d. T is **complete**.
- e. T is **consistent**.
- f. T is **categorical**.
- g. T is **equivalent** to T_2 ($T \equiv T_2$).

Fact B.3. *The following are well known facts:*

- a. If $T \vdash \phi$, then some finite subset of T proves ϕ (**compactness**).
- b. If T is complete and $T; \phi$ is consistent, then $T \vdash \phi$.
- c. If T is categorical, then T is complete.

Definition B.4. *More familiar notions:*

- a. T is **existential** iff it is equivalent to a set of prenex sentences, none of which have universal quantifiers.
- b. T is **universal** iff it is equivalent to a set of prenex sentences, none of which have existential quantifiers.
- c. T is **universal-existential** iff it is equivalent to a set of prenex sentences, each of which has all of its universal quantifiers preceding any of its existential quantifiers.
- d. ϕ is a **primitive formula** iff ϕ is an existential formula in prenex form whose matrix is a conjunction of atomic formulas and negations of atomic formulas.

Fact B.5. *Fairly familiar facts:*

- a. If T is existential, $\mathcal{A} \models T$, and $\mathcal{A} \subseteq \mathcal{B}$, then $\mathcal{B} \models T$.
- b. If T is universal and $\mathcal{A} \models T$, then any submodel of $\mathcal{A}$ also satisfies T .
- c. If T is universal-existential and $\mathcal{A}_i \models T$, for all $i > 0$, then $\bigcup \{\mathcal{A}_i\} \models T$.

C Model Completeness

This section presents the definition of *model completeness* and some basic results needed in Chapters 4 and 5.

There are several ways to show that a theory is complete. We use only one, which is based on A. Robinson's notion of *model completeness*.

Definition C.1. T is **model complete** iff T is consistent and for any two models, $\mathcal{A}$ and $\mathcal{B}$ of T , $\mathcal{A} \subseteq \mathcal{B}$ iff $\mathcal{A}$ is an elementary submodel of $\mathcal{B}$. [Monk, p. 355].

A model complete theory is not necessarily complete, unless the theory has a prime model.

Definition C.2. $\mathcal{A}$ is a **prime model** of T if $\mathcal{A} \models T$ and $\mathcal{A}$ can be embedded in any model of T . [Monk, p.359]

Fact C.3. If T is model complete and T has a prime model, then T is complete.

To show that a theory is model complete we rely, directly or indirectly, on a theorem of Monk's (see 4.2.1a), which is based on some equivalent formulations of model completeness:

Definition C.4.

- a. The **$\mathcal{A}$ -expansion** of L is the result of adding to L a constant for each element of $\mathcal{A}$.
- b. The **diagram** of $\mathcal{A}$ is the set of atomic sentences and negations of atomic sentences of the $\mathcal{A}$ -expansion of the language of $\mathcal{A}$ which are true in $\mathcal{A}$.

Fact C.5. The following are equivalent [Monk, p. 356]:

- a. T is model complete.
- b. For every model, $\mathcal{A}$, of T and every $\mathcal{A}$ -expansion L' of L , the $T \cup L'$ -diagram of L is complete.
- c. If $\mathcal{A}$ and $\mathcal{B}$ are models of T , $\mathcal{A} \subseteq \mathcal{B}$, ϕ is a universal formula, $x \in \mathcal{A}$, and $\mathcal{A} \models \phi(x)$, then $\mathcal{B} \models \phi(x)$.
- d. If $\mathcal{A}$ and $\mathcal{B}$ are models of T , $\mathcal{A} \subseteq \mathcal{B}$, ϕ is a primitive formula, $x \in \mathcal{A}$, and $\mathcal{B} \models \phi(x)$, then $\mathcal{A} \models \phi(x)$.

Bibliography

- [Bell] Bell and Slomson. *Models and Ultraproducts*. North-Holland Publishing Company, Amsterdam, 1971.
- [Cantor] Cantor, G. *Contributions to the Foundations of the Theory of Transfinite Numbers*. Dover Publications, New York, 1955.
- [Chang] Chang and Keisler. *Model Theory*. North-Holland Publishing Company, Amsterdam, 1973.
- [Dauben] Dauben, J.W. *Georg Cantor: His Mathematics and Philosophy of the Infinite*. Princeton University Press, Princeton, New Jersey, 1979.
- [Enderton] Enderton, H. *A Mathematical Introduction to Logic*. Academic Press, Inc., Orlando, Florida, 1972.
- [Frege] Frege, G. *The Foundations of Arithmetic*. Evanston, Illinois: Northwestern University Press, 1968.
- [Griffin] Griffin, H. *Elementary Theory of Numbers*. McGraw-Hill, New York, 1965.
- [Kurosh] Kurosh, A.G. *Lectures in General Algebra*. Pergamon Press, New York, 1965.
- [Monk] Monk, J. D. *Mathematical Logic* Springer-Verlag, New York, 1975.
- [Robinson] Robinson, A. and Zakon, E. *Transactions of the American Mathematical Society*. 96(1960), pp. 222-236.
- [Russell] Russell, B. *The Principles of Mathematics*. George Allen and Unwin, London, 1903. Second edition, reprinted by W.W. Norton and Company, New York, 1937.
- [Tarski] Tarski, A. *Undecidable Theories*. North-Holland Publishing Company, Amsterdam, 1968.

Index of Symbols

Logic

$\phi \wedge \psi$, 101
 $\phi \vee \psi$, 101
 $\phi \rightarrow \psi$, 101
 $\neg \phi$, 101
 $\phi \leftrightarrow \psi$, 101
 $\forall x \phi$, 101
 $\exists x \phi$, 101

Set Operations

$x \cup y$, 102
 $x \cap y$, 102
 $x - y$, 102
 $x; y$, 102
 $x \uplus y$, 102
 $\mathcal{P}(x)$, 102
 $|x|$, 102
 $x^{[k]}$, 102
 $\hat{x}^k$, 94

Arithmetic

$k \mid n$, 102
 $k!$, 102
 $[in + j]$, 102

Size ordering

$x \sim y$, 1
 $x < y$, 1
 $x \leq y$, 1
 $x > y$, 1
 $x \geq y$, 1

Model Theory

$T \vdash \phi$, 103
 $T \not\vdash \phi$, 103
 $T_1 \equiv T_2$, 103
 $\mathcal{A} \models \phi$, 103

$\mathcal{A} \not\models \phi$, 103

$\mathcal{A} \subseteq \mathcal{B}$, 103

Models and Domains

$\mathcal{A}$, 103
 $\hat{\mathcal{A}}$, 103
 $\mathcal{B}(\mathcal{A})$, 21
 $\mathcal{A}_n$, 86
 $\mathcal{A}_F$, 87
 $\hat{\mathcal{A}}$, 73
 $\mathcal{Q}$, 29
 $\mathcal{Q}_f$, 67
 $\mathcal{Q}_{f,n}$, 67
 $\mathcal{Q}_n$, 29
 $\mathcal{Q}_n$, 32
 QC , 27
 QC_n , 27
 $\mathbb{N}$, 25
 $\mathcal{S}_{\mathcal{A}}$, 45
 $\mathcal{Z}_{\mathcal{A}}$, 54
 $\hat{\mathcal{Z}}_{\mathcal{A}}$, 55

Sizing

$\tilde{x}$, 45
 $C(x)$, 31
 $\Delta_1(x)$, 31
 $\Delta_2(x)$, 31

$\rho(x)$, 32
 $\delta(x)$, 32
 $\theta(x)$, 32
 $\theta_f(x)$, 66
 $\alpha(x)$, 32
 $\beta(x)$, 32
 $\rho(x, y)$, 94
 $\sigma_{\mathcal{A}}[x]$, 45
 $\text{cvg}(x, y)$, 94

$\text{cvg}(x)$, 94

Node Construction

λ , 70

$\mathbf{L}(\lambda)$, 70

$\lambda \bullet m$, 70

$\mathbf{A}_\lambda$, 70

$\mathbf{a}_\lambda$, 70

$\hat{\mathcal{A}}$, 73

$\iota(\lambda)$, 75

$\mathbf{q}_\lambda$, 75

λ_n , 76

Index of Theories

Languages

L_C , 20

$L_{C<}$, 20

$L_{<}$, 20

L_N , 86

Predicates

$\text{Abstract}(x)$, 1

$\text{Atom}(x)$, 20

$\text{Div}_n(x)$, 37

$\text{Indisc}(x, y)$, 10

$\text{Mod}_{n,m}(x)$, 37

$\text{Sm}(x, y)$, 54

$\text{Sum}(x, y, z)$, 20

$\text{Unit}(x)$, 20

$\text{Times}_n(x, y)$, 37

Theories

ADIV_n , 37

ASSOC , 46

ASSOC2 , 54

$\text{ASYM}^<$, 10

ASYM^F , 10

$\text{ASYM}^>$, 10

ATLEAST_n , 22

BA , 22

BA_n , 22

BAI , 22

BASIC , 24

BDIV_J , 26

BDIV_n , 26

CA , 38

CAI , 40

CAI_f , 40

CORE , 13

CS , 21

CSI , 40

$\text{CANTOR}^<$, 9

DEF^+ , 16

DEF^+ , 24

$\text{DEF}^\sim$, 13

$\text{DEF}^>$, 10

$\text{DISJ}^\cup$, 24

DISJ^+ , 16

FUNC^+ , 16

DIV_J , 26

DIV_n , 25

EVEN , 9

EXACTLY_n , 22

EXCORE , 18

$\text{INDISC}^\sim$, 10

$\text{IRREF}^<$, 13

INF , 22

MAX , 46

MIN , 46

$\text{MOD}_{n,m}$, 25

MONOT , 16

ODD , 9

OUTPACING , 83

PCA , 47

PCS , 46

PSIZE , 46

QUASI , 10

$\text{REF}^\sim$, 10

$\text{REP}^<$, 9

RT_f , 40

SIZE , 24

$\text{SYM}^\sim$, 10

T_f , 40

$\text{TRANS}^\sim$, 10

$\text{TRANS}^<$, 10

$\text{TRANS}^>$, 10

TRICH , 13

UNIQ^- , 54

UNIQ^+ , 54

$\text{UNIQ}^\sim$, 46

Zg , 51

Zgm , 51

Subject Index

- $\mathcal{A}$ -finite set, 74
- $\mathcal{A}$ -near, 74
- abelian groups with identity, 51
- alternating pair, 85
- asymptotic density, 94
- basis, 21
- cancellable abelian semigroups with
 - identity, 51
- categorical, 104
- chain, 103
- charmed n -factor, 68
- class size, language of, 21
- classes, language of, 20
- common n -factor, 68
- compactness, 104
- complete, 103
- completion, 38
- congruence class, 27
- congruous, 40
- consistent, 103
- converges, 94
- density, 94
- depth, 70, 75
- diagram, 105
- diverges, 94
- elementary extension, 103
- equivalent, 104
- existential, 104
- existential assumption, 49
- expansion, 105
- extends, 70
- extension, 103
- filter, 87
- finite, 40
- finite class model, 14
- finite completion, 39
- finite intersection property, 87
- finite set model, 14
- functional assumptions, 49
- generated, 103
- incongruous, 40
- infinite completion, 39
- interpretation, 103
- isomorphic, 103
- isomorphic embedding, 103
- language of
 - class size ($L_{C<}$), 21
 - classes (L_C), 20
 - size ($L_{<}$), 21
 - subsets of $N(L_N)$, 86
- language of a theory, 103
- least common multiple, 36
- length, 70
- model complete, 104
- molecule, 64
- Monk mapping, 48
- Monk's Condition, 47
- n -congruence class, 27
- n -quasi-congruence class, 27
- N -semigroups, 52
- near, 27
- $(n.m)$ -partition, 68
- nodal embedding, 77
- node, 70
- node atom, 70

- node atoms, 71
- node set, 70
- node sets, 71

- one-one correspondence, 1
- outpaces, 83
- outpacing model, 84

- partial, 40
- PCA, 61
- prime model, 104
- primitive formula, 104
- principal, 87
- proves, 103

- quasi-congruence class, 27
- quasi-logical principles, 11
- quasi-nodal, 74

- reducible, 50
- remainder function, 40
- remainder theory, 40
- representation principle, 12
- roughly divisible, 25
- run, 91

- satisfaction, 103
- schematic function, 101
- shell, 70, 73
- simple order, 51
- simple translation, 48
- simply reducible, 50
- size, 30, 45, 66
- size model, 45, 46
- size, language of, 21
- solution, 40
- standard finite interpretation, 21
- standard interpretation, 21
- standard protomodel, 67
- submodel, 103
- subsets of $\mathbb{N}$, language of, 86

- τ -reducible, 50
- theory, 103
- theory of
 - abelian groups with identity, 51
 - cancellable abelian semigroups
 - with identity, 51
 - $\mathbb{N}$ -semigroups, 52
 - $\mathbb{Z}$ -groups, $\mathbb{Z}_g$, 51
 - $\mathbb{Z}$ -groups modulo I , 52
- total, 40
- translation, 48
- trichotomy, 12

- ultrafilter, 87
- uniformly τ -reducible, 50
- union of a chain, 103
- universal, 104
- universal-existential, 104

- $\mathbb{Z}$ -groups, $\mathbb{Z}_g$, 51
- $\mathbb{Z}$ -groups modulo I , 52
- 0-extends, 70